

# Molecular recognition of chloroform by divergent polymorphic transitions in *tert*-butylthiacalix[4]arene tetrasubstituted with *N*-(2-hydroxyethyl)carbamoylmethoxy groups in a lower rim

Goulnaz D. Safina, Olga M. Gavrilova, Marat A. Ziganshin,  
 Ivan I. Stoikov, Igor S. Antipin and Valery V. Gorbachuk\*

*A. M. Butlerov Institute of Chemistry, Kazan (Volga Region) Federal University, 420008 Kazan, Russian Federation.  
 Fax: +7 843 233 7416; e-mail: Valery.Gorbachuk@ksu.ru*

DOI: 10.1016/j.mencom.2011.09.022


The genuine molecular recognition of chloroform by a *tert*-butylthiacalix[4]arene derivative was observed, where the host remembers the guest above its elimination point showing two polymorphic transitions with opposite heat effects.

The polymorphism of solid receptors, giving the formation and collapse of metastable porous structures, can affect significantly their binding ability for vapours and gases.<sup>1–4</sup> In addition, it can be used for molecular recognition because the temperature and heat of polymorphic transition may depend on the receptor history, in particular, on the structure of the previously bound guest.<sup>2</sup>

Typically, a curve of differential scanning calorimetry (DSC) for clathrates of macrocyclic receptors has only *endo*-effects<sup>1,5,6</sup> or only *exo*-effects<sup>2,3</sup> corresponding to polymorphic transitions along with pseudopolymorphic transitions of guest release. Two different polymorphic transitions of a macrocyclic host with divergent heat effects in one run of thermoanalysis are very unusual, which had not been observed earlier. In this study, this phenomenon was observed for the clathrate of chloroform with *tert*-butylthiacalix[4]arene tetrasubstituted with *N*-(2-hydroxyethyl)carbamoylmethoxy groups in a lower rim, in the stable cone configuration (**1**),<sup>7</sup> above the point of the guest removal.

The combined method of thermogravimetry and DSC with the mass-spectrometric analysis of evolved gases (TG/DSC/MS experiment)<sup>3</sup> was used to analyze products of host **1** saturation by the vapours of methanol, ethanol, *n*-propanol and isopropanol, acetonitrile, propionitrile, benzene, toluene, pyridine, *n*-hexane, cyclohexane, dichloromethane, chloroform, tetrachloromethane, trichloroethylene and pentachloroethane with the relative vapour pressure  $P/P_0 = 1$  at  $T = 291$  K during 24 h<sup>†</sup> prepared as described elsewhere.<sup>3</sup> The TG/DSC/MS data are given in Figure 1.

In this experiment, thiacalixarene **1** was found to form relatively stable clathrates only with chloroform ( $1 \cdot 0.96 \text{CHCl}_3$ ) [Figure 1(a)], methanol ( $1 \cdot 0.98 \text{MeOH}$ ), ethanol ( $1 \cdot 0.70 \text{EtOH}$ ), *n*-propanol ( $1 \cdot 0.36 \text{PrOH}$ ), isopropanol ( $1 \cdot 0.25 \text{Pr}^i\text{OH}$ ), acetonitrile ( $1 \cdot 1.01 \text{MeCN}$ ) [Figure 1(d)], propionitrile ( $1 \cdot 0.43 \text{EtCN}$ ), dichloromethane ( $1 \cdot 0.58 \text{CH}_2\text{Cl}_2$ ) [Figure 1(e)], and pyridine ( $1 \cdot 0.41 \text{C}_5\text{H}_5\text{N}$ ) [Figure 1(f)]. Guest release from studied clathrates started at 30–40 °C, except for a clathrate of **1** with pyridine, which is stable to 90 °C. Ion thermograms (MS curve) show that only bound guests are evolved in this experiment. The other studied guests do not form stable clathrates under sample pre-equilibration conditions of thermal analysis: argon flow at room temperature for 20 min before heating.




The decomposition of the most of **1** clathrates with test guests shows only *endo*-effects of guest release in DSC curves [Figure 1(d)–(f)]. The exception is the DSC curve of  $1 \cdot 0.96 \text{CHCl}_3$  clathrate having an endothermic peak of chloroform removal at 50 °C, an exothermic peak at 116 °C with a heat value of  $\Delta H = -20.5 \text{ kJ mol}^{-1}$  and endothermic peak at 160 °C with  $\Delta H = 2.8 \text{ kJ mol}^{-1}$ . The mass loss of the sample in the temperature range of 105–125 °C is only 0.09% or 0.008 mol of  $\text{CHCl}_3$  per mole of **1**. An *exo*-peak in the DSC curve indicates a collapse of the loose  $\beta_0$ -phase formed during the decomposition of  $1 \cdot 0.96 \text{CHCl}_3$  clathrate ( $\beta$ -phase) and the formation of a dense form of the host ( $\alpha$ -phase).<sup>2,3</sup> Probably, a small residual amount of the guest in a  $\beta_0$ -phase stabilizes this metastable form of the host.<sup>4</sup> An *endo*-peak in the DSC curve without mass loss corresponds to a polymorphic transition from the dense  $\alpha$ -phase to a low-dense packing ( $\alpha'$ -phase).<sup>5,6</sup>

The loose  $\beta_0$ -phase of guest-free thiacalixarene **1** was prepared by the heating of  $1 \cdot 0.96 \text{CHCl}_3$  to 90 °C with a rate of 4 K min<sup>-1</sup> and fast cooling to 28 °C. Thermal analysis data for a  $\beta_0$ -phase sample are shown in Figure 1(b). According to the ion thermogram, this form of **1** does not contain chloroform. The enthalpies ( $-20.6$  and  $3.0 \text{ kJ mol}^{-1}$ ) and temperatures of polymorphic transitions for this sample coincide with the data observed for  $1 \cdot 0.96 \text{CHCl}_3$ .<sup>‡</sup>

The exothermic effect for host **1** is comparable with the collapse enthalpies of  $\beta_0$ -phase prepared from *tert*-butylcalix[6]arene<sup>2</sup> and adamantylcalix[4]arene.<sup>3</sup> Using a published procedure,<sup>2</sup> the free

<sup>†</sup> Thermoanalysis was performed in an argon atmosphere with a flow rate of 75 ml min<sup>-1</sup> at 4 K min<sup>-1</sup> for **1** clathrates and  $\beta_0$ -phase, and at 10 K min<sup>-1</sup> for  $\alpha'$ -phase of guest-free **1**. Clathrates were prepared by the saturation of host  $\alpha'$ -phase with guest vapours.

<sup>‡</sup> The saturation of all polymorphs of thiacalixarene **1** ( $\alpha$ -phase,  $\alpha'$ -phase and  $\beta_0$ -phase) by chloroform vapour with  $P/P_0 = 1$  at 298 K results in the formation of clathrate  $1 \cdot 0.96 \text{CHCl}_3$  with the same TG/DSC/MS curves as in Figure 1(a).



**Figure 1** Thermoanalysis data of (a) 1·0.96 CHCl<sub>3</sub> clathrate ( $\beta$ -phase), (b)  $\beta_0$ -phase of guest-free **1**, (c)  $\alpha'$ -phase of guest-free **1**, (d) 1·1.01 MeCN clathrate, (e) 1·0.58 CH<sub>2</sub>Cl<sub>2</sub> clathrate, (f) 1·0.41 C<sub>5</sub>H<sub>5</sub>N clathrate.


volume in  $\beta_0$ -phase of thiacalixarene **1** can be estimated by the ratio of host collapse enthalpies. This volume is more than a half of the space occupied by the guest in 1·0.96 CHCl<sub>3</sub> clathrate ( $\beta$ -phase). A similar calculation gives the free volume in  $\alpha'$ -phase of **1** being equal to 14% of the free volume in  $\beta_0$ -phase.

For comparison, the sample of thiacalixarene **1** ( $\alpha'$ -phase) without the guest was prepared by the heating of its clathrate with chloroform at 190 °C in a vacuum (100 Pa of argon atmosphere) for 6 h. Its TG/DSC analysis [Figure 1(c)] shows the absence of significant mass loss when heated from 30 to 300 °C. The most of the observed mass decrease in this range (about 1.3% by weight) is observed above 200 °C. The obtained DSC curve does not exhibit any polymorphic phase transitions. Above 320 °C, the decomposition of thiacalixarene **1** is observed.

The thermally stable  $\alpha'$ -phase and metastable  $\beta_0$ -phase prepared from clathrate **1** with chloroform were characterized using XRPD method.<sup>§</sup> The observed powder diffractograms (Figure 2) show differences in the positions and intensities of peaks, especially for  $2\theta$  values between 13 and 18°, which confirms the different crystal packing of the studied polymorphs. The presence of more pronounced peaks in the diffraction pattern (**1**) at large  $2\theta$  values indicates a higher degree of  $\alpha'$ -phase crystallinity.

The specific behaviour of thiacalixarene **1** upon heating may be due to the presence of groups in the lower rim of its macrocycle, which are capable to form intermolecular hydrogen bonds stabilizing a metastable loose  $\beta_0$ -phase upon guest removal and a loose  $\alpha'$ -phase upon further heating. Thus, host **1** remembers released chloroform, and only chloroform, allowing a genuine molecular recognition of this guest from others, including chloroalkanes.

<sup>§</sup> Powder X-ray diffraction (XRPD) data were collected using a Bruker D8 Advance diffractometer equipped with a Vario attachment and a Vantec linear position-sensitive detector. CuK $\alpha$  radiation (40 kV, 40 mA) and a graphite monochromator were used. The data were determined at room temperature in the reflection mode with a flat-plate sample. Patterns were recorded in the  $2\theta$  range between 5 and 50°, in 0.008° steps, with a step time of 1 s. The sample was lightly grounded and loaded into a standard sample holder.



**Figure 2** Powder XRD patterns of (1) thiacalixarene **1** ( $\alpha'$ -phase) after heating to 190 °C during 6 h and (2) porous metastable phase ( $\beta_0$ -phase) of thiacalixarene **1** prepared from clathrate 1·0.96 CHCl<sub>3</sub>.

This work was supported by programs of BRHE (REC007), Russian Foundation for Basic Research (grant no. 11-03-01215-a) and the Russian Ministry of Science and Education (state contract no. 14.740.11.0377). We are grateful to Dr. A. T. Gubaidullin (A. E. Arbuzov Institute of Organic and Physical Chemistry, Kazan Scientific Centre of the Russian Academy of Sciences) for performing XRPD experiments.

## References

- G. D. Safina, L. R. Validova, M. A. Ziganshin, I. I. Stoikov, I. S. Antipin and V. V. Gorbachuk, *Sens. Actuators, B: Chem.*, 2010, **148**, 264.
- A. V. Yakimov, M. A. Ziganshin, A. T. Gubaidullin and V. V. Gorbachuk, *Org. Biomol. Chem.*, 2008, **6**, 982.
- L. S. Yakimova, M. A. Ziganshin, V. A. Sidorov, V. V. Kovalev, E. A. Shokova, V. A. Tafeenko and V. V. Gorbachuk, *J. Phys. Chem. B*, 2008, **112**, 15569.
- P. K. Thallapally, S. J. Dalgarno and J. L. Atwood, *J. Am. Chem. Soc.*, 2006, **128**, 15060.
- M. A. Ziganshin, A. V. Yakimov, G. D. Safina, S. E. Solovieva, I. S. Antipin and V. V. Gorbachuk, *Org. Biomol. Chem.*, 2007, **5**, 1472.
- E. B. Brouwer, G. D. Enright, K. A. Udachin, S. Lang, K. J. Ooms, P. A. Halchuk and J. A. Ripmeester, *Chem. Commun.*, 2003, 1416.
- Y. Jin, C. Zhang, S. Gong and Y. Chen, *Synth. Commun.*, 2005, **35**, 3179.

Received: 29th April 2011; Com. 11/3719