

Selective reduction and dehydrogenation of 6-benzylideneoctahydropyrrolo[1,2-*a*]pyrimidines and 5-benzylidenehexahydropyrrolo[1,2-*a*]imidazoles as new approaches to *N*-(ω -aminoalkyl)pyrrolidines and bicyclic amidines

Konstantin N. Shavrin, Valentin D. Gvozdev and Oleg M. Nefedov

N-[3-(2-Benzyl-4,4-dimethylpyrrolidin-1-yl)propyl]-*N*-methylamine **2c** was obtained from amination **1c** in 83% yield. ¹H NMR, δ : 0.91 (s, 3H, CH₃), 1.04 (s, 3H, CH₃), 1.33 (dd, 1H, CHCHH₂CMe₂, ²J=12.7 Hz, ³J=8.4 Hz), 1.45 (dd, 1H, CHCHH₂CMe₂, ²J=12.7 Hz, ³J=7.2 Hz), 1.55-1.75 (m, 2H, NCH₂CH₂CH₂NH₂), 1.93 (d, 1H, NCHH₂CMe₂, J=9.1 Hz), 2.02 (br. s, 1H, NH), 2.16 (ddd, 1H, NCHH(CH₂)₂NHMe), ²J=11.7 Hz, ³J=7.2 Hz, ³J=5.1 Hz), 2.40 (dd, 1H, PhCHH, ²J=12.5 Hz, ³J=9.4 Hz), 2.41 (s, 3H, NHMe); 2.48-2.62 (m, 1H, PhCH₂CH), 2.61 (t, 2H, CH₂NHCH₃, J=6.7 Hz), 2.82-2.96 (m, 1H, NCHH(CH₂)₂NHCH₃), 2.92 (d, 1H, NCHH₂CMe₂, J=9 Hz), 3.00 (dd, 1H, PhCHH, J=12.5 Hz, J=3.6 Hz), 7.05-7.30 (m, 5H, Ph). ¹³C NMR, δ : 28.5 (NCH₂CH₂CH₂N); 29.1 (CH₃); 30.3 (CH₃); 35.6 (CMe₂); 36.5 (NCH₃); 41.1 (CH₂Ph); 46.3 (C-3, *cyclo*-C₄N); 51.1 (CH₂NMe); 53.2 (NCH₂); 66.9 (C-2, *cyclo*-C₄N); 68.3 (C-5, *cyclo*-C₄N); 125.9, 128.1, 129.2 (Ph); 140.0 (C-1, Ph). Mass spectrum, *m/z* (%): 169 ([M-CH₂Ph]⁺, 27), 112 ([M-CH₂Ph-C₃H₇N]⁺, 100), 44 ([C₂H₆N]⁺, 8). Found (%): C, 78.51; H, 10.60; N, 10.41. Calculated for C₁₇H₂₈N₂ (%): C, 78.41; H, 10.84; N, 10.76.

[3-(2-Benzylpyrrolidin-1-yl)propyl]amine **2e** was obtained from amination **1d** in 88% yield. ¹H NMR, δ : 1.35-1.75 (m, 8H, CHCH₂CH₂, NCH₂CH₂CH₂NH₂); 2.0-2.23 (m, 2H); 2.34-2.53 (m, 2H); 2.71 (t, 2H, CH₂NH₂, J=6.8 Hz); 2.80-3.03 (m, 2H); 3.09-3.20 (m, 1H). ¹³C NMR, δ : 21.7 (C-4, *cyclo*-C₄N); 30.2 (NCH₂CH₂CH₂N); 32.4 (C-3, *cyclo*-C₄N); 40.7, 40.9 (CH₂Ph, CH₂NH₂); 52.4 (NCH₂); 54.0 (C-5, *cyclo*-C₄N); 66.3 (C-2, *cyclo*-C₄N); 125.7, 128.0, 129.0 (Ph); 139.9 (C-1, Ph). Mass spectrum, *m/z* (%): 127 ([M-CH₂Ph]⁺, 42), 84 ([M-CH₂Ph-C₃H₇N]⁺). Found (%): C, 77.35; H, 9.95; N, 12.54. Calculated for C₁₄H₂₂N₂ (%): C, 77.01; H, 10.16; N, 12.83.

N-[3-(2-Benzylpyrrolidin-1-yl)propyl]-N-methylamine **2f** was obtained from aminal **1e** in 80% yield. ^1H NMR, δ : 1.40-1.78 (m, 7H, CHCH_2CH_2 , $\text{NCH}_2\text{CH}_2\text{CH}_2\text{NHCH}_3$); 2.05-2.28 (m, 2H); 2.35-2.55 (m, 2H); 2.41 (s, 3H, NCH_3); 2.62 (t, 2H, CH_2NHCH_3 , $J=6.8$ Hz); 2.85-3.05 (m, 2H); 3.15-3.26 (m, 1H). ^{13}C NMR, δ : 21.9 (C-4, *cyclo*- C_4N); 28.8 ($\text{NCH}_2\text{CH}_2\text{CH}_2\text{N}$); 30.3 (C-3, *cyclo*- C_4N); 36.5 (NCH_3); 40.9 (CH_2Ph); 50.8, 53.1 ($\text{NCH}_2\text{CH}_2\text{CH}_2\text{NCH}_3$); 54.1 (C-5, *cyclo*- C_4N); 66.4 (C-2, *cyclo*- C_4N); 125.8, 128.1, 129.1 (Ph); 139.9 (C-1, Ph). Mass spectrum, m/z (%): 141 ($[\text{M}-\text{CH}_2\text{Ph}]^+$, 31), 84 ($[\text{M}-\text{CH}_2\text{Ph}-\text{C}_3\text{H}_7\text{N}]^+$, 100), 44 ($[\text{C}_2\text{H}_6\text{N}]^+$, 10). Found (%): C, 77.59; H, 10.19; N, 12.25. Calculated for $\text{C}_{15}\text{H}_{24}\text{N}_2$ (%): C, 77.53; H, 10.41; N, 12.06.