

Effects of the electrostatic complexation between anionic pH-sensitive liposomes and star-shaped polycations on the release of the liposomal content

Andrey V. Sybachin, Olga V. Zaborova, Kristina M. Imelbaeva, Vyacheslav V. Samoshin, Vasily A. Migulin, Felix A. Plamper and Alexander A. Yaroslavov

Materials




Figure S1 The cationic PE star (schematic presentation).




Figure S2 Lipids (schematic presentation).

Experimental

A release of an encapsulated compound from PS¹⁻/EL/MOCH fliposomes was studied by conductometry using a CDM 83 Radiometer (Denmark) conductometer with a PP1042 platinum electrode.. The fliposomes loaded with NaCl solution were used. For this, during preparation the lipid film was dispersed in 1 M NaCl solution in 0.001 M Tris buffer with pH 7. After preparation of fliposomes the salt from the surrounding solution was eliminated by dialysis towards 0.001 M Tris buffer with pH 7. The dialysis process was repeated until conductivity of external solution became constant $42.9 \pm 0.2 \mu\text{S/cm}$. The conductivity of the 1 mg/mL fliposomes suspension in 0.001 M Tris buffer with pH 7 was $63.5 \pm 0.4 \mu\text{S/cm}$. A maximum conductivity after a complete destruction of fliposomes in the presence of an excess of a non-ionogenic Triton X-100 surfactant solution was measured $155.0 \pm 4.0 \mu\text{S/cm}$. The Triton X-100 by itself was found not to give impact in conductivity of the system.

The calibration curve demonstrating linear dependence of conductivity of Tris buffer upon concentration of added NaCl are presented on Figure S3.




Figure S3 Concentration dependence of conductivity of NaCl solution in 0.001 M Tris buffer with pH 7.

In the whole range of studied pH values the linear dependence of conductivity of system upon NaCl concentration. The calibration curve demonstrating linear dependence of conductivity of the most acidic acetate buffer pH 5 upon concentration of added NaCl are presented on Figure S4.

All measured conductance values were recalculated in relative conductivity by formula

$$\Omega = (\Omega_t - \Omega_0) / (\Omega_{\max} - \Omega_0),$$

where Ω_t is a current suspension conductivity, Ω_0 an initial conductivity and Ω_{\max} a maximum conductivity resulted after fliposomes disintegration by Triton X-100.




Figure S4 Concentration dependence of conductivity of NaCl solution in 0.001 M Acetate buffer with pH 5.