

Capturing polycyclic aromatic sulfur heterocycles in electron donor–acceptor complexes

Olga E. Eremina, Mariia V. Samodelova, Mariia V. Ferree, Tatyana N. Shekhovtsova and Irina A. Veselova

Table S1 Maximum wavelength of charge transfer complexes with DDQ in *n*-heptane, *n*-octane, and isoctane

with DDQ in CHCl ₃	λ_{\max} in <i>n</i> -heptane, nm	λ_{\max} in <i>n</i> -octane, nm	λ_{\max} in isoctane, nm
Dibenzothiophene	590 ± 10	590 ± 10	590 ± 10
4MDBT	605 ± 10	605 ± 10	620 ± 10
46DMDBT	610 ± 10	625 ± 10	620 ± 10
46DEDBT	615 ± 10	630 ± 10	640 ± 10

Table S2 Maximum wavelength of charge transfer complexes with TCNE in *n*-heptane, *n*-octane, and isoctane

with TCNE in CHCl ₃	λ_{\max} in <i>n</i> -heptane, nm	λ_{\max} in isoctane, nm
Dibenzothiophene	550 ± 10	550 ± 10
4MDBT	560 ± 10	565 ± 10
46DMDBT	560 ± 10	570 ± 10
46DEDBT	570 ± 10	580 ± 10




Figure S1 Bilogarithmic dependence of absorbance versus the π -acceptor concentration for CTCs: DBT:TCNE, 4MDBT:TCNE, 46DMDBT:TCNE, 46DEDBT:TCNE in isoctane.




Figure S2 Bilogarithmic dependence of absorbance versus the π -acceptor concentration for CTCs: DBT:TCNE, 4MDBT:TCNE, 46DMDBT:TCNE, 46DEDBT:TCNE in *n*-heptane.

The stoichiometric coefficients of the components of the complex correspond to the tangent of the slope of the curve ($tg\alpha$). The calculated stoichiometric ratios acceptor:donor are presented in Table S3.

Table S3 Calculation of stoichiometric of DBTs:TCNE charge transfer complexes in isoctane by Bent-French method

with TCNE	Equation	$tg\alpha$
Dibenzothiophene	$lgAbs = 1.18 \cdot lgc_A - 1.88$	1.18 ± 0.03
4MDBT	$lgAbs = 1.43 \cdot lgc_A - 2.18$	1.4 ± 0.1
46DMDBT	$lgAbs = 1.21 \cdot lgc_A - 1.89$	1.21 ± 0.06
46DEDBT	$lgAbs = 1.39 \cdot lgc_A - 2.10$	1.39 ± 0.09

Table S4 Calculation of stoichiometric of DBTs:TCNE charge transfer complexes in *n*-heptane by Bent-French method

with TCNE	Equation	$tg\alpha$
Dibenzothiophene	$lgAbs = 0.97 \cdot lgc_A - 1.88$	0.97 ± 0.03
4MDBT	$lgAbs = 1.32 \cdot lgc_A - 2.10$	1.3 ± 0.1
46DMDBT	$lgAbs = 1.41 \cdot lgc_A - 2.21$	1.4 ± 0.1
46DEDBT	$lgAbs = 1.21 \cdot lgc_A - 1.98$	1.2 ± 0.1




Figure S3 UV-vis spectra of individual solutions of DBT and its derivatives (10 mM) in chloroform.




Figure S4 UV-vis spectra of DBT, 4MDBT, 46DMDBT, and 46DEDBT (1 mM) in isoctane with DDQ (5 mM) in CHCl_3 .




Figure S5 UV-vis spectra of DBT, 4MDBT, 46DMDBT, and 46DEDBT (1 mM) in *n*-octane with different concentrations of DDQ in CHCl_3 .




Figure S6 UV-vis spectra of DBT, 4MDBT, 46DMDBT, and 46DEDDBT (1 mM) in *n*-heptane with different concentrations of DDQ in CHCl_3 .




Figure S7 UV-vis spectra of DBT, 4MDBT, 46DMDBT, and 46DEDDBT (1 mM) in isoctane with TCNE (5 mM) in CHCl_3 .




Figure S8 Linearized curves for calculation of stability constants of DBT, 4MDBT, 46DMDBT, and 46DEDBT in *n*-heptane.




Figure S9 Linearized curves for calculation of stability constants of DBT, 4MDBT, 46DMDBT, and 46DEDBT in *n*-octane.

Table S5 Stability constants (K_f) of charge transfer complexes with DDQ in *n*-heptane, *n*-octane, and isoctane

with DDQ in CHCl ₃	K_f in <i>n</i> -heptane, L/mol	K_f in <i>n</i> -octane, L/mol	K_f in isoctane, L/mol
Dibenzothiophene	20 ± 1	45 ± 5	20 ± 1
4MDBT	100 ± 12	30 ± 7	50.0 ± 0.3
46DMDBT	110 ± 13	58 ± 4	143 ± 8
46DEDBT	20 ± 3	20 ± 7	50.0 ± 0.3




Figure S10 Charge transfer complex of DBT with DDQ (A and B). LUMO of the charge transfer complex formed by DBT and DDQ (C).

Three-dimensional structures and orbital plots were produced with IQMol 2.15.