

## Synthesis and Structure of Phosphito- and Thiophosphatocavitands

Eduard E. Nifant'ev,<sup>\*a</sup> Vera I. Maslennikova,<sup>a</sup> Elena V. Panina,<sup>a</sup> Anna R. Bekker,<sup>a</sup> Larisa K. Vasyanina,<sup>a</sup> Konstantin A. Lyenko,<sup>b</sup> Mikhail Yu. Antipin<sup>c</sup> and Yurii T. Struchkov<sup>c</sup>

<sup>a</sup> V. I. Lenin Moscow Pedagogical State University, 119882 Moscow, Russian Federation. Fax: +7 095 248 0162

<sup>b</sup> Higher Chemical College, Russian Academy of Sciences, 125819 Moscow, Russian Federation.

<sup>c</sup> A. N. Nesmeyanov Institute of Organoelement Compounds, Russian Academy of Sciences, 117813 Moscow, Russian Federation. Fax: +7 095 135 5085


Phosphitocavitands, which on sulfurization result in all-*cis*-thiophosphatocavitands, have been obtained by phosphocyclization of octahydroxytetramethyl[1<sub>4</sub>] metacyclophane with amides of phosphorous acid, as confirmed by <sup>1</sup>H and <sup>31</sup>P NMR spectroscopy and X-ray analysis.

Cavitands are skeleton heterocycles that have a molecular cup structure. In recent years considerable study has been made of these compounds in terms of supramolecular chemistry.<sup>1,2</sup> However, the synthesis of their organophosphorus derivatives is little understood: only a short communication<sup>3</sup> and a patent<sup>4</sup> describing in general terms the possibility of metacyclophane phosphocyclization by dichlorides of trivalent phosphorus acids have been published.

The present paper is concerned with the synthesis of evidence for a possible reaction of metacyclophanes with phosphocavitands by cyclophosphorylation of octahydroxy-[1<sub>4</sub>]metacyclophanes with phosphorous amides, commonly used in fine organic synthesis to create complex skeleton systems in particular.<sup>5</sup> Furthermore, some structural and other fundamental problems of cavitand chemistry are solved. Octahydroxytetramethyl(tetraphenyl)[1<sub>4</sub>]metacyclophanes **1** and both phosphorous triamides **2a,b** and phosphorous diamidoester **2c** are used as starting substances (Scheme 1).<sup>†</sup>

Phosphorylation of tetramethyl derivatives **1a** proceeds selectively and results in the 1,3,2-dioxaphosphocine systems **3a–c**. According to <sup>31</sup>P NMR spectroscopy, the crude products comprise individual compounds with minor amounts of stereoisomers. On reprecipitation, the minor isomers are separated or transformed into the major ones. Tetraphenyl-metacyclophane **1b**, because of its conformational features,<sup>6</sup> is phosphorylated with more difficulty. Therefore, we failed to separate stereoisomers resulting from cyclophosphorylation<sup>7</sup> and to isolate pure cavitands. This fact demonstrates the severity of the synthetic problem under investigation.


<sup>†</sup> Cyclophosphorylation was performed in dioxane at 65–100 °C under vigorous stirring. Cavitands **3a–c** were isolated by reprecipitation from dioxane on adding hexane. Yields of the products **3a–c** were 70–80%, m.p. 270–275 °C (decomp.). Their identity and structure were proved by means of <sup>1</sup>H and <sup>31</sup>P NMR spectroscopy (Table 1). The elemental analysis data are consistent with the theoretical values.



Scheme 1

This work begins to study the chemical features of the phosphocavitands obtained. It is found that they are alkylated with difficulty and that the phosphoamides **3a,b** are not subject to alcoholysis under the usual conditions. Thus, the compounds obtained manifest notable chemical peculiarities; for example, they add sulfur to form thiophosphoryl derivatives **4**<sup>†</sup> (Scheme 2).<sup>§</sup>

It is significant that the cavitand molecule adds sulfur stereoselectively only through the axial orbitals of trivalent



Scheme 2

phosphorus. Such a stereoconsistent sulfurization phenomenon for a polyphosphite system appears to be the first to be revealed. The axially-oriented sulfur atoms seem to make

**Table 1** <sup>1</sup>H and <sup>31</sup>P NMR spectral parameters of phosphocavitands.

Cavitands	$\delta_P^a$ (ppm)	$\delta_H$ /ppm <sup>b</sup> ( <sup>3</sup> J/Hz)					
		H <sub>m</sub>	H <sub>o</sub>	CHCH <sub>3</sub>	CHCH <sub>3</sub>	NCH <sub>2</sub> CH <sub>3</sub>	NCH <sub>2</sub> CH <sub>3</sub>
<b>3a<sup>c</sup></b>	141.3	7.26	6.49	4.80	1.73 (7.3 HH)		
<b>3b</b>	142.6	7.27	6.49	4.80	1.73 (7.4 HH)	3.28 (9.8 HP)	1.17
<b>3c<sup>d</sup></b>	130.1	7.39	6.61	4.83	1.79 (8.3 HH)		
<b>4a</b>	69.2	7.21	6.55	4.75	1.77 (7.3 HH)		
<b>4b<sup>e</sup></b>	66.7	7.38	6.56	4.74	1.84 (7.4 HH)	3.43 (13.2 HP)	1.20

<sup>a</sup> 32.4 MHz, CHCl<sub>3</sub>, against H<sub>3</sub>PO<sub>4</sub>. <sup>b</sup> 400 MHz, CDCl<sub>3</sub>, against TMS. <sup>c</sup> NMe: 2.8 (<sup>3</sup>J<sub>HP</sub> 10.3). <sup>d</sup> CD<sub>2</sub>Cl<sub>2</sub>, OMe: 3.90 (<sup>3</sup>J<sub>HP</sub> 8.8). <sup>e</sup> NMe: 2.95 (<sup>3</sup>J<sub>HP</sub> 12.4).

<sup>†</sup> It should be noted that a paper on phosphocyclization of octahydroxy[1<sub>4</sub>]metacyclophane **1** by dichlorophosphate has recently been published.<sup>5</sup> In that case the reaction occurred with no selectivity and resulted in a mixture of diastereoisomers.


<sup>§</sup> Sulfur addition occurs in dioxane on heating at 50–60 °C for 0.5 h. Yields of cavitands **4a,b** are 90–95%, m.p. > 300 °C.

the cavitand bowl deeper and create possibilities for the building of new rims within it.

The structural peculiarities of the cavitands **4a,b** were studied by NMR spectroscopy (see Table 1) and those of the compound **4b** by X-ray analysis.<sup>†</sup>

The crystals of **4b** contain a molecule of cavitand, a solvate molecule of methanol and chloroform. Within a crystal of tetragonal symmetry, the molecule of **4b** occupies a special position and possesses fourth-order crystallographic symmetry. The fourth-order axis passes through the macrocyclic nucleus normally to the central plane of the molecule. The geometric characteristics for the structure of **4b** have the expected values for an all-*cis*-cavitand (see Fig. 1). The solvate molecule of methanol resides in the cavity (on the axis 4 of the cavitand) and forms a hydrogen bond O–H...S with thiophosphoryl groups (the distance S...O 3.19 Å fits the hydrogen bonds of moderate strength).

The work was performed within the program 'Russian Foundation for Basic Research'.



**Fig. 1** Principal bond lengths (Å) and bond angles (°) in the structure **4b**: P=S 1.904 (2), P–O(1) 1.610(4), P–O(2) 1.607(4), P–N 1.644(4); S–P–N 117.3(2), S–P–O(1) 115.6(2), S–P–O(2) 114.8(1), O(1)–P–O(2) 102.8(2), N–P–O(1) 100.1(2), N–P–O(2) 104.1(2).

<sup>†</sup> *Crystallographic data for 4b*: Tetragonal crystals, space group *P4/n*, at –90 °C, *a* = *b* = 18.883(5) Å, *c* = 10.233(3) Å, *V* = 3649(2) Å<sup>3</sup>, *Z* = 4, *d*<sub>calc</sub> = 1.441 g cm<sup>–3</sup>,  $\mu(\text{MoK}\alpha) = 7.09 \text{ cm}^{-1}$ , *F*(000) = 1628. Intensities of 5225 reflections were measured on a Syntex-P2<sub>1</sub> diffractometer at –90 °C (MoK $\alpha$  radiation,  $\Theta/2\Theta$  scan,  $2\Theta < 50^\circ$ ), and 2255 independent observed ones with  $I > 2\sigma(I)$  were used in calculations and refinement. The structure was solved by a direct method and refined by least-squares in an anisotropic-isotropic (H atoms and atoms of the methyl alcohol molecule) approximation to *R* = 6.74%, *w*<sub>R</sub> = 7.40% and GOF = 1.54. All calculations were performed using the program SHELXTL PLUS on an IBM PC/AT computer. Atomic coordinates, thermal parameters, bond lengths and bond angles have been deposited at the Cambridge Crystallographic Data Centre (CCOC), see 'Notice to Authors', *Mendelev Commun.*, 1995, issue no. 1.

## References

- 1 D. J. Cram, *Angew. Chem.*, 1988, **100**, 1041.
- 2 D. J. Cram, *Nature*, 1992, **356**, 29.
- 3 W. Xu, P. Rourke, J. J. Vittal and R. J. Puddephatt, *J. Chem. Soc., Chem. Commun.*, 1993, 145.
- 4 M. Foa and S. Strologo, *Eur. Patent*, 487036, 1992 (*Chem. Abstr.*, 117: P 213761v).
- 5 E. E. Nifantsev and M. K. Gratchev, *Usp. Khim.*, 1994, **63**, 602 (*Russ. Chem. Rev.*, 1994, **63**, 575).
- 6 A. G. Högberg, *J. Am. Chem. Soc.*, 1980, **102**, 6046.
- 7 L. Abis, E. Dalkanale, A. Du vosel and S. Spera, *J. Chem. Soc., Perkin Trans. 2*, 1990, 2075.
- 8 T. Lippmann, E. Dalkanale, and G. Mann, *Tetrahedron Lett.*, 1994, **35**, 1685.

Received: Moscow, 14th February 1995  
Cambridge, 6th April 1995; Com. 5/01016K