

Nitrodeiodination of Polyiodopyrazoles: a Convenient Synthesis of 4-Nitroiodopyrazoles


Eugene V. Tretyakov and Sergei F. Vasilevsky*

Institute of Chemical Kinetics and Combustion, Siberian Branch of the Russian Academy of Sciences, 630090 Novosibirsk, Russian Federation. Fax: +7 3832 35 2350; e-mail: vasilev@kinetics.nsk.su

A number of 3,4-, 4,5-diiodo- and 3,4,5-triiodo-1-methylpyrazoles **2a–f** have been converted into the corresponding almost inaccessible 3-, 5-iodo- and 3,5-diiodo-1-methyl-4-nitropyrazoles **3a–f** by nitration with an HNO₃–H₂SO₄ mixture.

Intramolecular cyclization of functionally substituted vicinal aromatic acetylenic compounds has recently become increasingly important as a method for the synthesis of heterocyclic condensed systems.¹ The use of pyrazoles containing an iodine atom and functional groups (NO₂, NH₂) could provide a route to inaccessible bicyclic compounds which bear aromatic or non-aromatic and pyrazole rings. Furthermore, this could give additional information on the reasons for the different reaction pathways in the heterocyclization of functionally substituted acetylenic derivatives of benzene and pyrazole.^{2–4} Only vicinal iodonitropyrazoles form such functionally substituted acetylenic pyrazoles, because on the one hand, only iodo- (but not chloro- or bromo-) derivatives of pyrazoles react with the acetylenic moiety, and on the other hand, the nitro group may be readily converted into amino and then diazo groups. The fact that the oxidative iodination of 1-methylpyrazoles using I₂–HIO₃ is an efficient method allows us to prepare some iodo-derivatives containing even electron-withdrawing substituents in the ring.^{5–7} However, attempts to iodinate the pyrazole ring with a nitro, amino or acetylamino group in the 4-position proved unsuccessful.⁵ A general synthetic approach to iodo-4-nitro-1-methylpyrazoles is still unknown.

In the present communication we report a convenient synthetic procedure for the preparation of the iodonitropyrazoles by means of exhaustive iodination of a pyrazole ring[†] with subsequent nitration (Scheme 1).




Scheme 1

We examined 4,5-diiodo-1,3- and 3,4-diiodo-1,5-dimethylpyrazoles **2a** and **2b**, respectively, in order to determine the general applicability of *ipso* nitration. The replacement of the iodine atom proceeded smoothly at 50 °C in a solution of concentrated (or fuming) nitric acid in 94% sulfuric acid in

[†] In a typical experiment, 0.01 mol of **1a–f**, 2.03 g (0.008 mol) of I₂ and 0.7 g (0.004 mol) of HIO₃ in a mixture of 40 ml of AcOH, 4 ml of 30% H₂SO₄ and 5 ml of CCl₄ were heated at 80–85 °C for 2.5 h (**1a–d,f**) or 40 h (**1e**) until the iodine colour disappeared, after which the mixture was diluted with 150 ml of water, the precipitate was filtered and washed with a saturated solution of Na₂SO₃ and with water. Recrystallisation of the residue from ethanol–water gave diiodo-derivatives **2a–d,f**. 3,4,5-Triiodo-1-methylpyrazole **2e** was synthesised in a similar manner except for the fact that we used 3 equiv. of iodinating mixture, 3.05 g (0.012 mol) of I₂ and 1.06 g (0.006 mol) of HIO₃.⁵

yields of 90–93%.[‡] As to the literature data, Grimmett⁸ and Stepanov^{9,10} reported on the nitrodebromination of 4-bromo-1-methylpyrazoles, while the nitrodeiodination of 4-iodo-1-methylpyrazoles as reported has failed.¹¹

Determination of the nitro group position in **3a** was based on both the difference in the reactivity towards electrophiles of the 4- and other positions of the ring, and a comparison of the physical and spectroscopic data of **3a** with the known 5-iodo-1,3-dimethyl-4-nitropyrazole.³ In addition, the reduction of **3a** gave 4-amino-5-iodo-1,3-dimethylpyrazole³ rather than 5-amino-4-iodo-1,3-dimethylpyrazole.⁵ Iodonitropyrazole **3b** was also obtained by nitrodecarboxylation of 3-iodo-1,5-dimethylpyrazole-4-carboxylic acid **4**,¹² which unambiguously proved its structure (Scheme 2).[§]



Scheme 2

As expected, the nitration of 3,4,5-triiodo-1-methylpyrazole **2f** requires more time. The yield of 3,5-diiodo-1-methyl-4-nitropyrazole **3f** was rather good (74%).

Taking into account the fact that 4-chloro- and 4-bromo-1-methylpyrazolecarboxylic acids failed to undergo *ipso* nitration^{9,11} it is interesting to clarify the possibility of nitrodeiodination in the presence of a carboxylic group. We have found that both 3,4-diiodo- **2c** and 4,5-diiodopyrazolecarboxylic acid **2d** give iodo-4-nitropyrazolecarboxylic acids **3c,d** in 51–68% yields.

In order to establish the generality of nitrodeiodination and its applicability to the synthesis of iodonitropyrazoles, we nitrated 4,5-diiodo-1-methyl-3-nitropyrazole **2e**. It has been demonstrated that even the vicinal nitro group did not prevent *ipso* substitution of the iodine atom in the 4-position of pyrazole. Moreover, surprisingly, TLC testing of the reaction mixture after it had been heated at 50 °C for 10 min showed the absence of starting material. Work-up gave 5-iodo-3,4-dinitro-1-methylpyrazole **3e** in 94% yield. The reasons for the acceleration of nitrodeiodination in this case

[‡] General method for the preparation of iodo-4-nitropyrazoles **3a–f**. A mixture of fuming nitric acid (10 ml) and 94% sulfuric acid (10 ml) was added dropwise to 7.76 mmol diiodopyrazole **2a–f** in 94% sulfuric acid (30 ml). The mixture was heated at 50 °C for the period of time indicated in Table 1 (TLC), then poured into ice (70–80 g). The cold (0–5 °C) suspension was filtered out and the solid was washed with a saturated solution of Na₂SO₃ and with water. Recrystallisation of the residue gave 4-nitroiodopyrazoles **3a–f**. The results are summarised in Tables 1 and 2.

[§] A mixture of fuming nitric acid (13.5 ml) and 98% sulfuric acid (13.5 ml) was added dropwise to 8 g (30 mmol) **4** in 98% sulfuric acid (4.5 ml). The mixture was heated at 70–80 °C for 1 h, then poured into ice. The cold (0–5 °C) suspension was filtered and the solid washed with water. Recrystallisation of the residue from EtOH gave 4.7 g (58.7%) of the 4-nitroiodopyrazole **3b**.

Table 1 Reaction times, yields and melting points of 4-nitroiodopyrazoles 3a–f.

No	Time/min	Yield (%)	M.p./°C (crystallisation solvent)	Formula
3a	40	92.7	149–151 ^a (EtOH–H ₂ O 10:1)	–
3b	40	90.3	134–135 (CHCl ₃)	C ₅ H ₆ N ₃ O ₂ I
3c	180	51.5	194–194.5 (C ₆ H ₆ –CHCl ₃ 1:1)	C ₅ H ₄ N ₃ O ₄ I
3d	180	68.4	224–225 (decomp.) (C ₆ H ₆ –CHCl ₃ 1:1)	C ₅ H ₄ N ₃ O ₄ I
3e	10	93.6	77–78 (hexane–C ₆ H ₆ 3:1)	C ₄ H ₃ N ₄ O ₄ I
3f	90	74.2	226–226.5 (decomp.) (EtOH)	C ₄ H ₃ N ₃ O ₂ I ₂

^a Ref. 3: 150–151 °C (tetrachloromethane).

as well as for an extension of the synthesis by the above route are currently being investigated.

In all cases studied only one product was isolated. This presents an excellent route to iodo-4-nitropyrazoles, which are useful synthons.

We thank Professor M. S. Shvartsberg for useful discussions. This work was supported by grant no. 95-03-08928a, awarded by the Russian Foundation for Basic Research, to whom the authors express their gratitude.

References

- 1 M. S. Shvartsberg, *Izv. Sib. Otd. Akad. Nauk SSSR, Ser. Khim. Nauk*, 1983, **4**, 96 (in Russian).
- 2 S. F. Vasilevskii and E. V. Tretyakov, *Abstracts of Papers*, 14th International Congress of Heterocyclic Chemistry, University of Antwerp RUCA, Antwerp, Belgium, 1993, Abstract PO3-165.
- 3 S. F. Vasilevskii, T. V. Anisimova and M. S. Shvartsberg, *Izv. Akad. Nauk SSSR, Ser. Khim.*, 1983, 688 (*Bull. Acad. Sci. USSR, Div. Chem. Sci.*, 1983, **32**, 626).
- 4 S. F. Vasilevskii, *Izv. Sib. Otd. Akad. Nauk SSSR, Ser. Khim. Nauk*, 1986, **4**, 105 (in Russian).
- 5 S. F. Vasilevskii and M. S. Shvartsberg, *Izv. Akad. Nauk SSSR, Ser. Khim.*, 1980, 1071 (*Bull. Acad. Sci. USSR, Div. Chem. Sci.*, 1980, **29**, 778).

Table 2 Analytical and characteristic IR spectral data.

No	Calc./Found (%)			IR (CHCl ₃): ν _{max} /cm ⁻¹
	C	H	I	
3b	20.47	2.33	47.46	1332, 1555
	20.49	2.26	47.52	
3c	20.22	1.36	42.73	1310, 1330, 1410, 1540, 1760, 3540
	20.26	1.31	42.71	
3d	20.22	1.36	42.73	1320, 1330, 1405, 1540, 1770, 3530
	20.30	1.23	42.70	
3e	16.12	1.01	42.58	1330, 1360, 1560, 1570
	16.16	1.09	42.60	
3f	12.68	0.80	66.99	1330, 1500
	12.84	0.73	66.87	

- 6 S. F. Vasilevskii, A. L. Belov and M. S. Shvartsberg, *Izv. Sib. Otd. Akad. Nauk SSSR, Ser. Khim. Nauk*, 1985, **5**, 100 (in Russian).
- 7 J. H. Boyer, in *Organic Nitro Chemistry. Nitroazoles. The C-Nitro Derivatives of Five-membered N- and N,O-Heterocycles*, VCH, Deerfield Beach, 1986, p. 186.
- 8 K. Chang, M. R. Grimmett, D. D. Ward and R. T. Weavers, *Aust. J. Chem.*, 1979, **32**, 1727.
- 9 V. P. Perevalov, M. A. Andreeva, L. I. Baryshnenkova, Y. A. Manaev, G. S. Yamburg, B. I. Stepanov and V. A. Dubrovskaya, *Khim. Geterotsikl. Soedin.*, 1983, 1676 [*Chem. Heterocycl. Compd. (Engl. Transl.)*, 1983, 1326].
- 10 V. P. Perevalov, L. I. Baryshnenkova, M. A. Andreeva, Y. A. Manaev, I. A. Denisova, B. I. Stepanov and V. I. Seraya, *Khim. Geterotsikl. Soedin.*, 1983, 1672 [*Chem. Heterocycl. Compd. (Engl. Transl.)*, 1983, 1322].
- 11 Y. A. Manaev, M. A. Andreeva, V. P. Perevalov, B. I. Stepanov, V. A. Dubrovskaya and V. I. Seraya, *Zh. Obshch. Khim.*, 1982, **52**, 2592 [*J. Gen. Chem. USSR (Engl. Transl.)*, 1982, 2291].
- 12 S. F. Vasilevskii, V. A. Gerasimov and M. S. Shvartsberg, *Izv. Akad. Nauk SSSR, Ser. Khim.*, 1981, 902 (*Bull. Acad. Sci. USSR, Div. Chem. Sci.*, 1981, **30**, 683).

Received: Moscow, 26th June 1995
Cambridge, 14th August 1995; Com. 5/04294A