

Aminaleacetal of dimethylaminomethylenemalonaldehyde – a novel reagent for the synthesis of substituted aminopolyenes

Zhanna A. Krasnaya,* Yuliya V. Smirnova, Galina V. Krystal and Valentin S. Bogdanov

N. D. Zelinsky Institute of Organic Chemistry, Russian Academy of Sciences, 117913 Moscow, Russian Federation.

Fax: +7 095 135 5328


Synthesis of the previously unknown bis-aminaleacetal of dimethylaminomethylenemalonaldehyde has been developed, condensation of which with malonates affords trienic δ -dimethylaminotetraesters.

As shown previously, amins and aminaleacetals of conjugated β -dimethylaminoaldehydes $\text{Me}_2\text{N}(\text{CH}=\text{CH})_n\text{CH}(\text{Y})\text{NMe}_2$ ($n = 1-4$, $\text{Y} = \text{OMe}$, NMe_2) are convenient reagents for the synthesis of various types of aminopolyenes, because they condense easily with ketones, β -diketones and CH-acids, on reactive methyl and methylene groups, without the need for catalysts.¹⁻⁴

Investigations of aminopolyenes have shown that they possess a number of specific properties, *viz.* solvatochromism, thermochromism, fluorescence, laser-emitted light generation, *etc.*⁵⁻⁷

In this work a method of synthesizing the previously unknown bis-aminaleacetal of dimethylaminomethylenemalonaldehyde has been developed, with the aim of obtaining new types of substituted aminopolyenes. Bis-aminaleacetal **1** was prepared from bis-perchlorate **3** (Scheme 1).[†]

The synthesis of starting compound **3** has been described earlier.⁸ However, our attempts to obtain bis-perchlorate **3** by formylation of chloroacetic acid **2a** according to this procedure⁸ were unsuccessful. Only inorganic salts were isolated from the reaction mixture following the recommended treatment with ice and a concentrated aqueous solution of



Scheme 1

NaClO_4 .


Bis-perchlorate **3** was, however, prepared after changing the order of adding the reagents [POCl_3 was added dropwise to a solution of **2a** in absolute dimethylformamide (DMF)] and on treatment of the reaction solution with EtOH, H_2O and 35% aqueous HClO_4 . The yield of bis-perchlorate **3**, however, was not greater than 25%.

Formylation of bromoacetic acid **2b** as above allowed us to increase the yield of salt **3** (up to 76%).

Bis-aminaleacetal **1** was prepared by interaction of bis-perchlorate **3** with MeOK in benzene solution in the presence of NHMe_2 . From ^1H NMR spectral data compound **1** was shown to be bis-aminaleacetal (NMe_2 and OMe group ratio 1:1). The arrangement of NMe_2 and OMe (Scheme 1) is conditional, since OMe groups can be located on one carbon atom, while NMe_2 groups are also on another. Following

[†] **Compound 3.** To a stirred mixture of **2b** (20.8 g, 0.15 mol) in dry DMF (65 ml) was added dropwise POCl_3 (49 ml, 0.54 mol) over 3 h at 5–12°C. The mixture was heated slowly and stirred at 90°C for 10 h. The reaction solution was then cooled to –20 to –25°C and treated dropwise with EtOH (60 ml), H_2O (120 ml) and 35% aqueous HClO_4 (70 ml) with stirring. The temperature of the reaction mixture was kept at –20 to –25°C for 2 h; the resulting solid was filtered off, washed with EtOH and dried under reduced pressure over CaCl_2 at room temperature to give 43.7 g (76%) of bis-perchlorate **3**. Characteristic data for compound **3** are given in ref. 8.

Compound 1. To a stirred suspension of MeOK in dry benzene, that had been obtained from K (4.9 g, 0.13 mol) and dry MeOH (5.3 ml, 0.13 mol) in a nitrogen atmosphere at 6°C was added bis-perchlorate **3** (12 g, 0.03 mol) and a solution of Me_2NH in dry benzene (1:1, 12 ml). The mixture was then heated rapidly to 25°C and stirred for 1 h. The resulting solution was filtered, solids washed extensively with dry C_6H_6 and the solvent removed under reduced pressure. The oil was distilled *in vacuo* to give 5 g (60%) of bis-aminaleacetal of dimethylaminomethylenemalonaldehyde **1**, bp 68–70°C (0.6 mmHg); n_D^{20} 1.4788; ^1H NMR (C_6D_6): 6.12 and 6.20 (1H, CH=), 4.50, 4.42 and 4.28 (2H, CH), 3.25–3.38 (6H, m, OMe), 2.75 (6H, m, NMe_2), 2.46–2.56 (12H, m, NMe_2).



Scheme 2

distillation *in vacuo* (0.6 mmHg), bis-aminaleacetal **1** was found to possess satisfactory storage stability at $-5-0^{\circ}\text{C}$.

We found that the yield of **1** depends, to a great extent, on the molar ratio of MeOK to bis-perchlorate **3** (2:1 30%; 4:1 60%; 6:1 94%).

Condensation of bis-aminaleacetal **1** with malonates **4a,b**, without catalysts, at room temperature affords novel trienic tetraesters **5a,b** (Scheme 2).[‡]

These reactions have shown, in principle, the possibility of involving both aminaleacetal groups in the condensation of bis-aminaleacetal **1**.

The results of studies on the condensation of bis-aminaleacetal **1** with various ketones and CH-acids will be the subject of our future publications.

[‡] *General procedure.* To a solution of **1** (0.5 g, 1.9 mmol) in dry benzene (2 ml) at room temperature was added malonate **4a** (0.43 ml, 3.8 mmol). The reaction mixture was allowed to stand for 24 h at room temperature, then the solvent was removed under reduced pressure, absolute ether was added to the oil residue and the mixture cooled to -70°C . The resulting yellow powder was filtered off, washed with absolute ether to give 0.37 g **5a** (30%), mp $109-113^{\circ}\text{C}$ (from MeOH).

Spectroscopic data. Compound **5a**: $^1\text{H NMR}$ (CDCl_3): 7.52 (2H, s, H- β), 6.99 (1H, s, H- δ), 3.77 (6H, s, COOMe), 3.74 (6H, s, COOMe), 3.16 (6H, s, NMe₂); UV $\lambda_{\text{max}}/\text{nm}$ (EtOH): 372 (ϵ 32600).

Compound **5b**: yield 31%; mp $92-94^{\circ}\text{C}$; $^1\text{H NMR}$ (CDCl_3): 7.50 (2H, s, H- β), 6.95 (1H, s, H- δ), 4.22 (4H, q, CH₂), 4.19 (4H, q, CH₂), 3.14 (6H, s, NMe₂), 1.27 (6H, t, Me), 1.26 (6H, t, Me); UV $\lambda_{\text{max}}/\text{nm}$ (EtOH): 372 (ϵ 21800).

References

- 1 Zh. A. Krasnaya, T. S. Stytsenko, V. A. Petukhov and V. F. Kucherov, *Izv. Akad. Nauk SSSR, Ser. Khim.*, 1976, 595 (*Bull. Acad. Sci. USSR, Div. Chem. Sci.*, 1976, **25**, 577).
- 2 Zh. A. Krasnaya and T. S. Stytsenko, *Izv. Akad. Nauk SSSR, Ser. Khim.*, 1981, 1060 (*Bull. Acad. Sci. USSR, Div. Chem. Sci.*, 1981, **30**, 820).
- 3 Zh. A. Krasnaya, E. V. Grechkina and V. S. Bogdanov, *Izv. Akad. Nauk, Ser. Khim.*, 1993, 1235 (*Russ. Chem. Bull.*, 1993, **42**, 1179); 1994, 285 (*Russ. Chem. Bull.*, 1994, **43**, 264).
- 4 Zh. A. Krasnaya, Yu. V. Smirnova, A. S. Tatikolov, V. S. Bogdanov, E. V. Nikishova and V. L. Savel'ev, *Izv. Akad. Nauk, Ser. Khim.*, 1995, 537 (*Russ. Chem. Bull.*, 1995, **44**, 520).
- 5 Zh. A. Krasnaya, T. S. Stytsenko, B. M. Uzhinov, S. A. Krashakov and V. S. Bogdanov, *Izv. Akad. Nauk SSSR, Ser. Khim.*, 1983, 2084 (*Bull. Acad. Sci. USSR, Div. Chem. Sci.*, 1983, **32**, 1882).
- 6 L. A. Shvedova, A. S. Tatikolov, A. P. Darmanyan, V. A. Kuz'min and Zh. A. Krasnaya, *Dokl. Akad. Nauk SSSR*, 1984, **276**, 164 [*Dokl. Phys. Chem. (Engl. Transl.)*, 1984, **276**, 398].
- 7 L. A. Shvedova, A. S. Tatikolov, V. A. Kuz'min, Zh. A. Krasnaya and A. R. Bekker, *Dokl. Akad. Nauk SSSR*, 1984, **276**, 654 [*Dokl. Phys. Chem. (Engl. Transl.)*, 1984, **276**, 457].
- 8 Z. Arnold, *Collect. Czech. Chem. Commun.*, 1965, **30**, 2125.

Received: Moscow, 28th June 1995

Cambridge, 2nd November 1995; Com. 5/04296H