

# Stereoselective synthesis of triterpene 3-*O*-2-deoxy- $\alpha$ -glycosides


Lidiya A. Baltina, Oksana B. Flekhter\* and Ekaterina V. Vasiljjeva

Institute of Organic Chemistry, Ufa Research Centre of the Russian Academy of Sciences, 450054 Ufa, Russian Federation.  
Fax: +7 3472 356 066

A synthesis of 3-*O*-2-deoxy- $\alpha$ -glycosides of glycyrrhetic acid and allobetulin was carried out by electrophilic glycosylation of triterpene alcohols with D-glucal acetate in the presence of (*sym*-collidin)<sub>2</sub>IClO<sub>4</sub>.

A great number plants, of various species, and sea invertebrates contain triterpene glycosides (TG). Natural TGs are known by their high and diverse level of biological activity.<sup>1</sup> A synthesis of TG modelling different natural analogs (*ginseng* and *Glycyrrhizae glabra* glycosides)<sup>2–4</sup> has attracted a great deal of attention over the last few years. An electrophilic glycosylation of alcohols with readily available glycols is a worthwhile method of preparing glycosides.<sup>5–8</sup>

the basis of literature data for 2-deoxy- $\alpha$ -glycosides<sup>8</sup> and *via* a comparison with the spectra of initial compounds. <sup>13</sup>C NMR spectra of aglycone fragments of glycosides **3–5** are similar to those for initial triterpenes.<sup>13–15</sup> In the <sup>13</sup>C NMR spectra of glycosides **3a–c**† signals due to carbinol atoms at C3 were observed at 84.3–84.7 ppm. On going from genins **2a–c** to glycosides **3a–c** the C3 signal was shifted to a lower field (5.3–6.7 ppm). Signals of hydrocarbon atoms at C2' in the



**Scheme 1** Reagents and conditions: i, (*sym*-collidin)<sub>2</sub>IClO<sub>4</sub>, CH<sub>2</sub>Cl<sub>2</sub>, 4A molecular sieves; ii, 10% Pd/C, ethyl acetate; iii, 5% KOH–MeOH.

Deoxyglycosides are structural units of many major biologically active materials (antibiotics, deoxyribonucleotides, heart glycosides, *etc.*). A synthesis of triterpene 2-deoxyglycosides has not previously been described in the literature.

We have carried out a synthesis of the oleanane-type 3-*O*-2-deoxy- $\alpha$ -glycosides **5a–c** *via* electrophilic glycosylation of triterpene alcohols with D-glucal acetate<sup>9</sup> **1** in the presence of iodonium di(*sym*-collidin) perchlorate<sup>6</sup> (Scheme 1). 18 $\beta$ -Glycyrrhetic acid<sup>10</sup> being one of the main biologically active triterpenoids of licorice roots (*Glycyrrhizae glabra* L. and *G. uralensis* F.), its 11-deoxo-analog, as methyl esters **2a,b**, and allobetulin<sup>11</sup> **2c** from the triterpene fraction of bark (*Betula pendula*) were used as the alcohol components in the synthesis reported here.

The reaction of triterpenoids **2a–c** with glycol **1** proceeds stereoselectively in methylene dichloride at room temperature. A mole ratio of reagents were used in the presence of 4A molecular sieves and (*sym*-collidin)<sub>2</sub>IClO<sub>4</sub> for 4–5 h to give 2-deoxy-2-iodo- $\alpha$ -D-manno-glycosides **3a–c** in 70–80% yield, which were separated by column chromatography on silica gel. Hydride deiodination of the protected 2-iodo- $\alpha$ -glycosides **3a–c** in the presence of 10% Pd/C in ethyl acetate<sup>8</sup> and deacetylation by 5% KOH in methanol<sup>12</sup> led to the triterpene-free 3-*O*-2-deoxy- $\alpha$ -D-arabino-hexopyranosides **5a–c** in yields of 75–80%. Elemental analyses and theoretical computations for all the product compounds are in a good agreement with the proposed structures. The glycoside structure was assigned by <sup>1</sup>H and <sup>13</sup>C NMR spectroscopy. Signal determination in the spectra of freshly prepared glycosides was carried out on

† **3a**: C<sub>43</sub>H<sub>63</sub>O<sub>11</sub> *M*<sub>r</sub> = 882.9, amorph; [ $\alpha$ ]<sub>D</sub><sup>20</sup> + 82° (*c* 0.03, CHCl<sub>3</sub>); UV (MeOH),  $\lambda_{\text{max}}$ /nm: 248.2 (lg *e* 3.59); <sup>13</sup>C NMR (CDCl<sub>3</sub>,  $\delta$ , ppm): 22.4 (C2), 84.6 (C3), 200.5 (C11), 128.5 (C12), 169.5 (C13), 177.1 (C30), 51.9 (C31), 98.5 (C1'), 30.9 (C2'), 69.3 (C3'), 67.9 (C4'), 69.7 (C5'), 62.4 (C6'); <sup>1</sup>H NMR (CDCl<sub>3</sub>,  $\delta$ , ppm): 0.73, 0.79, 0.94, 1.05, 1.07, 1.22 (s, 7CH<sub>3</sub>), 1.10–1.97 (m, CH<sub>2</sub>, CH), 1.99, 2.02, 2.03 (s, 3Ac), 2.25 (s, 1H, H9), 2.77 (d, 1H, H18, *J* = 13.7 Hz), 3.15 (dd, 1H, H3, *J*<sub>3,2e</sub> = 4.6 Hz, *J*<sub>3,2a</sub> = 11.1 Hz), 3.64 (s, 3H, OCH<sub>3</sub>), 4.00–4.12 (m, 1H, H5', 2H, H6'), 4.41 (dd, 1H, H2', *J*<sub>2',1'</sub> = 1.1, *J*<sub>2',3'</sub> = 4.2 Hz), 4.56 (dd, 1H, H3', *J*<sub>3',2'</sub> = 4.2, *J*<sub>3',4'</sub> = 9.5 Hz), 5.23 (d, 1H, H1', *J*<sub>1',2'</sub> = 1.1 Hz), 5.29 (t, 1H, H4', *J*<sub>4',3'</sub> = *J*<sub>4',5'</sub> = 9.5 Hz), 5.59 (s, 1H, H12).

**3b**: C<sub>43</sub>H<sub>65</sub>O<sub>10</sub> *M*<sub>r</sub> = 868.9, amorph; [ $\alpha$ ]<sub>D</sub><sup>20</sup> + 75° (*c* 0.02, CHCl<sub>3</sub>); <sup>13</sup>C NMR (CDCl<sub>3</sub>,  $\delta$ , ppm): 22.3 (C2), 84.7 (C3), 29.7 (C11), 122.4 (C12), 144.4 (C13), 177.8 (C30), 51.7 (C31), 98.4 (C1'), 31.0 (C2'), 69.3 (C3'), 67.7 (C4'), 69.6 (C5'), 62.3 (C6'); <sup>1</sup>H NMR (CDCl<sub>3</sub>,  $\delta$ , ppm): 0.75, 0.82, 0.94, 1.10 (s, 7CH<sub>3</sub>), 1.10–2.00 (m, CH<sub>2</sub>, CH), 2.05, 2.06, 2.09 (s, 3Ac), 3.20 (dd, 1H, H3, *J*<sub>3,2e</sub> = 4.6 Hz, *J*<sub>3,2a</sub> = 11.1 Hz), 3.63 (s, 3H, OCH<sub>3</sub>), 4.15–4.25 (m, 1H, H5', 2H, H6'), 4.47 (dd, 1H, H2', *J*<sub>2',1'</sub> = 1.0 Hz, *J*<sub>2',3'</sub> = 4.1 Hz), 4.61 (dd, 1H, H3', *J*<sub>3',2'</sub> = 4.1 Hz, *J*<sub>3',4'</sub> = 9.4 Hz), 5.25 (s, 1H, H12), 5.31 (d, 1H, H1', *J*<sub>1',2'</sub> = 1.0 Hz), 5.37 (t, 1H, H4', *J*<sub>4',3'</sub> = *J*<sub>4',5'</sub> = 9.5 Hz).

**3c**: C<sub>42</sub>H<sub>65</sub>O<sub>9</sub> *M*<sub>r</sub> = 840.9, amorph; [ $\alpha$ ]<sub>D</sub><sup>20</sup> + 54° (*c* 0.04, CHCl<sub>3</sub>); <sup>13</sup>C NMR (CDCl<sub>3</sub>,  $\delta$ , ppm): 22.4 (C2), 84.3 (C3), 26.2 (C12), 87.8 (C19), 71.2 (C28), 28.8 (C30), 98.3 (C1'), 31.0 (C2'), 69.2 (C3'), 67.6 (C4'), 69.5 (C5'), 62.3 (C6'); <sup>1</sup>H NMR (CDCl<sub>3</sub>,  $\delta$ , ppm): 0.79, 0.82, 0.86, 0.92, 0.98, 1.01 (s, 7CH<sub>3</sub>), 1.00–2.00 (m, CH<sub>2</sub>, CH), 2.03, 2.05, 2.07 (s, 3Ac), 3.18–3.23 (br. signal, 1H, H3), 3.42 (d, 1H, H28, *J* = 7.9 Hz), 3.50 (s, 1H, H19), 3.76 (d, 1H, H28, *J* = 7.9 Hz), 4.17–4.28 (m, 1H, H5', 2H, H6'), 4.47 (dd, 1H, H2', *J*<sub>2',1'</sub> = 1.0 Hz, *J*<sub>2',3'</sub> = 3.9 Hz), 4.62 (dd, 1H, H3', *J*<sub>3',2'</sub> = 3.9 Hz, *J*<sub>3',4'</sub> = 9.3 Hz), 5.30 (d, 1H, H1', *J*<sub>1',2'</sub> = 1.0 Hz), 5.35 (t, 1H, H4', *J*<sub>4',3'</sub> = *J*<sub>4',5'</sub> = 9.3 Hz).

spectra of 2-iodoglycosides **3a–c** were observed at 30.9–31.0 ppm, in the spectra of protected 2-deoxyglycosides **4a–c**<sup>‡</sup> at 35.1–35.7 ppm and in the spectra of free glycosides **5a–c**<sup>§</sup> at 38.1–38.6 ppm. Anomeric hydrogen atoms at C1' of the carbohydrate residues in the spectra of **3a–c** resonate at 98.3–98.5 ppm and at 93.4–93.5 ppm in the spectra of glycosides **5a–c**,<sup>13–15</sup> which provides evidence for the existence of an  $\alpha$ -glycosidic linkage.<sup>16</sup> The  $\alpha$ -configuration of the glycosidic linkage was confirmed by a doublet of anomeric proton H1' in the <sup>1</sup>H NMR spectra at 5.27 ppm with a spin–spin coupling constant (SSCC) of  $J_{H1',H2'} = 1.1$  Hz. The H2' proton signal was observed at 4.4 ppm as a doublet of doublets with SSCCs of  $J_{H2',H1'} = 1.1$  Hz and  $J_{H2',H3'} = 4.0$  Hz. The small value of SSCC for both H1' and H2' protons demonstrates their diequatorial position and, hence, the  $\alpha$ -glycosidic linkage and axial position of iodine atom. In the <sup>13</sup>C NMR spectra in the regime NOE of **3a–c**, the geminal constant value  $^2J_{C1',H2'}$  is small (<1 Hz), which confirms an axial position of the aglycone.<sup>17</sup> Thus, glycosides **3a–c** were found to be of  $\alpha$ -D-manno-configuration and in a <sup>4</sup>C<sub>1</sub>(D) conformation. The formation of  $\alpha$ -D-manno-glycosides has been observed during the glycosylation of steroid alcohols by glycals<sup>8</sup> and a synthesis of disaccharides.<sup>6</sup>

<sup>‡</sup> **4a**: C<sub>43</sub>H<sub>64</sub>O<sub>11</sub>  $M_r = 757.0$ , amorph;  $[\alpha]_D^{20} + 87^\circ$  ( $c$  0.04, CHCl<sub>3</sub>); UV (MeOH),  $\lambda_{max}/nm$ : 248.2 ( $\lg \epsilon$  3.66); <sup>13</sup>C NMR (CDCl<sub>3</sub>,  $\delta$ , ppm): 21.8 (C2), 87.2 (C3), 93.2 (C1'), 35.1 (C2'), 69.3 (C3'), 68.3 (C4'), 69.7 (C5'), 62.5 (C6').

**4b**: C<sub>43</sub>H<sub>66</sub>O<sub>10</sub>  $M_r = 743.0$ , amorph;  $[\alpha]_D^{20} + 78^\circ$  ( $c$  0.04, CHCl<sub>3</sub>); <sup>13</sup>C NMR (CDCl<sub>3</sub>,  $\delta$ , ppm): 21.9 (C2), 83.0, (C3), 93.2 (C1'), 35.7 (C2'), 69.3 (C3'), 68.4 (C4'), 69.7 (C5'), 62.6 (C6').

**4c**: C<sub>42</sub>H<sub>66</sub>O<sub>9</sub>  $M_r = 715.0$ , amorph;  $[\alpha]_D^{20} + 56^\circ$  ( $c$  0.07, CHCl<sub>3</sub>); <sup>13</sup>C NMR (CDCl<sub>3</sub>,  $\delta$ , ppm): 21.9 (C2), 82.8 (C3), 93.2 (C1'), 35.6 (C2'), 69.3 (C3'), 68.3 (C4'), 69.7 (C5'), 62.5 (C6').

<sup>§</sup> **5a**: C<sub>37</sub>H<sub>58</sub>O<sub>8</sub>  $M_r = 630.9$ , decomp. 210–212 °C;  $[\alpha]_D^{20} + 95^\circ$  ( $c$  0.02, CHCl<sub>3</sub>); UV (MeOH),  $\lambda_{max}/nm$ : 248.0 ( $\lg \epsilon$  3.81); <sup>13</sup>C NMR (CDCl<sub>3</sub>,  $\delta$ , ppm): 21.7 (C2), 81.7 (C3), 93.4 (C1'), 38.1 (C2'), 72.0 (C3'), 69.2 (C4'), 72.5 (C5'), 62.2 (C6').

**5b**: C<sub>37</sub>H<sub>60</sub>O<sub>7</sub>  $M_r = 616.9$ , decomp. 214–216 °C;  $[\alpha]_D^{20} + 83^\circ$  ( $c$  0.05, CHCl<sub>3</sub>); <sup>13</sup>C NMR (CDCl<sub>3</sub>,  $\delta$ , ppm): 21.8 (C2), 82.0 (C3), 93.5 (C1'), 38.3 (C2'), 71.9 (C3'), 69.3 (C3'), 69.3 (C4'), 72.5 (C5'), 62.1 (C6').

**5c**: C<sub>36</sub>H<sub>60</sub>O<sub>6</sub>  $M_r = 588.9$ , decomp. 202–204 °C;  $[\alpha]_D^{20} + 61^\circ$  ( $c$  0.06, CHCl<sub>3</sub>); <sup>13</sup>C NMR (CDCl<sub>3</sub>,  $\delta$ , ppm): 21.9 (C2), 82.0 (C3), 93.5 (C1'), 38.6 (C2'), 71.9 (C3'), 69.3 (C4'), 72.8 (C5'), 62.5 (C6').

## References

- 1 Th. Schople and K. Hiller, *Pharmazie*, 1990, **45**, 313.
- 2 L. N. Atopkina, N. F. Samoshina and N. I. Uvarova, *Khim. Prir. Soedin.*, 1989, **6**, 813 [*Chem. Nat. Compd. (Engl. Transl.)*, 1989, **6**, 690].
- 3 S. Saito, K. Kuroda, Y. Hayashi, Y. Sasaki, Y. Nagamara, K. Nishida and I. Ishiguro, *Chem. Pharm. Bull.*, 1991, **39**, 2333.
- 4 S. Saito, S. Sumita, Y. Kanda and Y. Sasaki, *Chem. Pharm. Bull.*, 1994, **42**, 1016.
- 5 A. G. Tolstikov and G. A. Tolstikov, *Usp. Khim.*, 1993, **62**, 621 (*Russ. Chem. Rev.*, 1993, **62**, 579).
- 6 R. W. Friesen and S. J. Danishefsky, *J. Am. Chem. Soc.*, 1989, **111**, 6656.
- 7 R. L. Halcomb and S. J. Danishefsky, *J. Am. Chem. Soc.*, 1989, **111**, 6661.
- 8 J. Thiem, S. Kopper and J. Schwentner, *Liebigs Ann. Chem.*, 1985, **11**, 2135.
- 9 W. Roth and W. Pigman, *Methods Carbohydr. Chem.*, 1963, **2**, 405.
- 10 G. A. Tolstikov and M. I. Goryaev, *Glitsirretovaya kislota (Glycyrrhetic acid)*, Nauka, Alma-Ata, 1966, p. 72 (in Russian).
- 11 F. N. Lugemwa, F.-Y. Huang, M. D. Bentley, M. J. Mendel and A. R. Alford, *J. Agric. Food Chem.*, 1990, **38**, 49.
- 12 M. Hirooka, N. Morishima, E. Kaji, Y. Mori and Sh. Zen, *Yakugaku Zasshi*, 1989, **109**, 544.
- 13 G. S. Ricca, B. Danieli, G. Palmisano, H. Duddeck and M. H. A. Elgamal, *Org. Magn. Reson.*, 1978, **11**, 163.
- 14 G. A. Tolstikov, L. M. Khalilov, L. A. Baltina, R. M. Kondratenko, A. A. Panasenko and E. V. Vasil'eva, *Khim. Prir. Soedin.*, 1985, **5**, 645 [*Chem. Nat. Compd. (Engl. Transl.)*, 1985, **5**, 605].
- 15 L. E. Odinokova, G. I. Oshitok, V. A. Denisenko, V. Ph. Anuphriev, A. M. Tolkach and N. I. Uvarova, *Khim. Prir. Soedin.*, 1984, **2**, 182 [*Chem. Nat. Compd. (Engl. Transl.)*, 1984, **2**, 168].
- 16 K. Tori, S. Seo, Y. Yoshimura, H. Arita and Y. Tomita, *Tetrahedron Lett.*, 1977, 179.
- 17 J. L. Marshall, *Methods in Stereochemical Analysis*, Chemie International, Florida, 1983, vol. 2, p. 30.

Received: Moscow, 3rd July 1995

Cambridge, 14th August 1995; Com. 5/04610F