

# The first example of synthesis of aluminacyclopropanes catalysed by $(\eta^5\text{-C}_5\text{H}_5)_2\text{TiCl}_2$

Usein M. Dzhemilev,\* Askhat G. Ibragimov, Leila O. Khafizova, Sergey V. Rusakov and Leonard M. Khalilov


Institute of Petrochemistry and Catalysis, the Bashkortostan Republic Academy of Sciences, 450075 Ufa, Russian Federation.  
Fax: +7 3472 31 2750; e-mail: root@ink.bashkiria.su

The synthesis of a series of cyclic organoaluminium compounds, aluminacyclopropanes, which have never previously been described, has been carried out by  $\alpha$ -olefin cyclometallation with  $\text{EtAlCl}_2$  in the presence of a chlorine ion acceptor, metallic magnesium and  $\text{Cp}_2\text{TiCl}_2$  ( $\text{Cp} = \eta^5\text{-C}_5\text{H}_5$ ) catalyst.

Recently we have developed regio- and stereoselective methods to synthesize aluminacyclopropanes,<sup>1,7</sup> aluminacyclopentanes,<sup>8,9</sup> and 1,4-dialuminium compounds<sup>10</sup> using zirconium catalysts. The reactions were assumed to proceed in the presence of zirconium-containing metallacycles as key intermediates.

To extend the study on the catalytic cyclometallation of olefins by alkylalanes to five-member organoaluminium compounds (OAC) and to understand the possibilities of applying this approach to the synthesis of aluminacyclopropanes we have studied the cyclometallation of  $\alpha$ -olefins with  $\text{EtAlCl}_2$ -Mg reagent in the presence of  $\text{Cp}_2\text{TiCl}_2$  catalyst.


The catalyst for cycloaluminum of  $\alpha$ -olefins to cyclic OAC were chosen in the expectation that titana- **1** or zirconacyclopropane **2** intermediates<sup>11-15</sup> generated *in situ* would be transmetalated with  $\text{EtAlCl}_2$  to give substituted aluminacyclopropanes **3**.



To confirm the supposition proposed, we have studied the cyclometallation of styrene by  $\text{EtAlCl}_2$ -Mg effected by  $\text{Cp}_2\text{ZrCl}_2$  and  $\text{Cp}_2\text{TiCl}_2$  with high selectivity and catalytic activity in reactions of hydro-, carbo- and cyclo-metallization of olefins.<sup>3,16-18</sup>


As a result we have found that  $\text{Cp}_2\text{ZrCl}_2$  (under the chosen conditions) promoted the generation of aluminacyclopentanes,<sup>5</sup> whereas  $\text{Cp}_2\text{TiCl}_2$  directed the reaction towards formation of aluminacyclopropanes. Thus, 1-ethyl-2-phenylaluminacyclopropane **4** was formed in the interaction of styrene with equimolar amounts of  $\text{EtAlCl}_2$  and Mg (powder) in the presence of catalytic amounts of  $\text{Cp}_2\text{TiCl}_2$  (3–5 mol%) in THF for 6–8 h at room temperature (22–23 °C) in 65–85% yield. The latter was isolated by distillation under reduced pressure in an inert atmosphere. The product was colourless, readily decomposed in air and was stable in an inert atmosphere. Each experiment gave 5–10% of 1-ethyl-2,4-diphenylaluminacyclopentane together with **4**.<sup>†</sup>

Ethylbenzene **5** was formed in the hydrolysis of the reaction mixture by 5% HCl, and 1,2-dideuterioethylbenzene **6** was formed in deuteriolysis, the latter being identified by spectral methods. The structure of **4** was confirmed by NMR spectroscopy.<sup>‡</sup> The <sup>13</sup>C NMR spectrum of **4** contains two pairs of broadened signals in the high field region. A wide triplet signal at 1.80 ppm and a quartet signal at 9.20 ppm might be assigned



to the ethyl group on the aluminium atom. Methylene and methyne carbon atoms of the aluminacyclopropane fragment had an intense broadened triplet at 1.50 ppm and a doublet at 9.92 ppm.

According to the literature data and our experimental results we suppose that titanacyclopropane **7** was formed initially in the cycloaluminum reaction under the chosen conditions and that it was then transmetalated with  $\text{EtAlCl}_2$  to give **4**. Subsequent incorporation of a styrene molecule into **7** with respect to the Ti-C bond gave titanacyclopentane **8**. Transmetalation of the latter with the initial  $\text{EtAlCl}_2$  was found to promote formation of a minor (5–10%) product **9**.




<sup>‡</sup> Spectral data for 1-ethyl-2-phenylaluminacyclopropane **4**: <sup>13</sup>C NMR ( $\text{C}_6\text{D}_6$ , 22.5 MHz, Jeol FX 90Q)  $\delta$ : 1.50 (br t,  $\text{C}^3$ ), 1.80 (br t,  $\text{C}^4$ ), 9.20 (q,  $\text{C}^5$ ), 9.92 (br d,  $\text{C}^2$ ), 126.44 (d,  $\text{C}^9$ ), 128.53 (d,  $\text{C}^8$ ,  $\text{C}^{10}$ ), 129.12 (d,  $\text{C}^7$ ,  $\text{C}^{11}$ ), 137.73 (s,  $\text{C}^6$ ).

For 1,2-dideuterioethylbenzene **6**: <sup>1</sup>H NMR ( $\text{CDCl}_3$ , 300 MHz, Bruker AM300)  $\delta$ : 1.16 (d,  $\text{CH}_2\text{D}$ ,  $^3J_{\text{H-H}}$  7.6 Hz), 2.57 (t,  $\text{CHD}$ ,  $^3J_{\text{H-H}}$  7.6 Hz), 7.02–7.24 (m, 5H,  $\text{C}_6\text{H}_5$ ). <sup>13</sup>C NMR ( $\text{CDCl}_3$ , 75 MHz, Bruker AM300)  $\delta$ : 15.35 (t,  $\text{C}^1$ ,  $J_{\text{C-D}}$  22.0 Hz), 28.57 (t,  $\text{C}^2$ ,  $J_{\text{C-D}}$  22.0 Hz), 125.65 ( $\text{C}^6$ ), 127.90 ( $\text{C}^4$ ,  $\text{C}^8$ ), 128.37 ( $\text{C}^5$ ,  $\text{C}^7$ ), 144.23 ( $\text{C}^3$ ).

<sup>†</sup> A 50 ml flask equipped with a magnetic stirrer under dry argon at 0 °C was charged with styrene (1.04 g, 10 mmol), THF (10 ml), Mg powder (0.288 g, 12 mmol),  $\text{Cp}_2\text{TiCl}_2$  (0.10 g, 0.5 mmol) and  $\text{EtAlCl}_2$  (1.524 g, 12 mmol), and the temperature was increased to room temperature (22–23 °C). The reaction mixture was stirred for 8 h, then treated with 5% HCl or  $\text{DCI}/\text{D}_2\text{O}$ .

To extend the field of application of styrene cycloaluminum to **4** under the chosen conditions (5 mol% of  $\text{Cp}_2\text{TiCl}_2$ , THF, 22–23 °C, 8 h) we carried out cycloaluminum of *p*-methylstyrene and 1,4-diphenylbuta-1,3-diene with  $\text{EtAlCl}_2$  to give aluminacyclopropanes **10** and **11** in 80% yield.



The structures of **10** and **11** were confirmed by spectral methods and deuteriolysis of the latter to hydrocarbons **12** and **13**.<sup>§</sup>

Cycloaluminum of alkyl-, cycloalkyl- and Si- and Sn-containing  $\alpha$ -olefins effected by  $\text{Cp}_2\text{TiCl}_2$  will be discussed in future publications.

The research work was supported by the Russian Foundation for Basic Research (grant nos. 96-15-97288, 97-03-33131 and 95-03-09807).

<sup>§</sup> Spectral data for **10**:  $^{13}\text{C}$  NMR ( $\text{C}_6\text{D}_6$ , 22.5 MHz, Jeol FX 90Q)  $\delta$ : 10.03 (br. d,  $\text{C}^2$ ), 2.63 (br. t,  $\text{C}^3$ ), 1.59 (br. t,  $\text{C}^4$ ), 9.84 (q,  $\text{C}^5$ ), 135.24 (s,  $\text{C}^6$ ), 128.06 (d,  $\text{C}^7$ ), 128.52 (d,  $\text{C}^8$ ), 136.30 (s,  $\text{C}^9$ ), 21.03 (q,  $\text{C}^{10}$ ).

For **12**:  $^{13}\text{C}$  NMR ( $\text{CDCl}_3$ , 75 MHz, Bruker AM300)  $\delta$ : 15.52 (t,  $\text{C}^1$ ,  $J_{\text{C-D}} = 19.05$  Hz), 28.13 (t,  $\text{C}^2$ ,  $J_{\text{C-D}} = 19.05$  Hz), 141.22 (s,  $\text{C}^3$ ), 127.82 (d,  $\text{C}^4$ ), 129.06 (d,  $\text{C}^5$ ), 135.04 (s,  $\text{C}^6$ ), 21.04 (q,  $\text{C}^7$ ).

For **13**:  $^{13}\text{C}$  NMR ( $\text{CDCl}_3$ , 75 MHz, Bruker AM300)  $\delta$ : 35.63 (t,  $\text{C}^1$ ,  $J_{\text{C-D}} = 19.60$  Hz), 34.55 (t,  $\text{C}^2$ ,  $J_{\text{C-D}} = 19.60$  Hz), 129.71 (d,  $\text{C}^3$ ), 126.88 (d,  $\text{C}^4$ ), 137.59 (s,  $\text{C}^5$ ), 130.34 (d,  $\text{C}^6$ ), 128.41 (d,  $\text{C}^7$ ), 125.84 (d,  $\text{C}^8$ ), 141.55 (s,  $\text{C}^9$ ), 128.41 (d,  $\text{C}^{10}$ ,  $\text{C}^{11}$ ), 125.96 (d,  $\text{C}^{12}$ ).

## References

- U. M. Dzhemilev, A. G. Ibragimov, A. P. Zolotarev, R. R. Muslukhov and G. A. Tolstikov, *Izv. Akad. Nauk SSSR, Ser. Khim.*, 1989, 207 (*Bull. Acad. Sci. USSR, Div. Chem. Sci.*, 1989, **38**, 194).
- U. M. Dzhemilev, A. G. Ibragimov, A. P. Zolotarev, R. R. Muslukhov and G. A. Tolstikov, *Izv. Akad. Nauk SSSR, Ser. Khim.*, 1990, 2831 (*Bull. Acad. Sci. USSR, Div. Chem. Sci.*, 1990, **39**, 2570).
- U. M. Dzhemilev, *Tetrahedron*, 1995, **51**, 4333.
- U. M. Dzhemilev, A. G. Ibragimov, A. B. Morozov, L. M. Khalilov, R. R. Muslukhov and G. A. Tolstikov, *Izv. Akad. Nauk SSSR, Ser. Khim.*, 1991, 1141 (*Bull. Acad. Sci. USSR, Div. Chem. Sci.*, 1991, **40**, 1022).
- U. M. Dzhemilev, A. G. Ibragimov, A. B. Morozov, R. R. Muslukhov and G. A. Tolstikov, *Izv. Akad. Nauk SSSR, Ser. Khim.*, 1991, 1607 (*Bull. Acad. Sci. USSR, Div. Chem. Sci.*, 1991, **40**, 1425).
- U. M. Dzhemilev, A. G. Ibragimov and A. B. Morozov, *Mendeleev Commun.*, 1992, 26.
- U. M. Dzhemilev, A. G. Ibragimov, A. P. Zolotarev, L. M. Khalilov and R. R. Muslukhov, *Izv. Akad. Nauk, Ser. Khim.*, 1992, 386 (*Bull. Russ. Acad. Sci., Div. Chem. Sci.*, 1992, **41**, 300).
- U. M. Dzhemilev and A. G. Ibragimov, *J. Organometal. Chem.*, 1994, **466**, 1.
- U. M. Dzhemilev, A. G. Ibragimov and A. P. Zolotarev, *Mendeleev Commun.*, 1992, 135.
- U. M. Dzhemilev, A. G. Ibragimov, M. N. Arhgaliev and R. R. Muslukhov, *Izv. Akad. Nauk, Ser. Khim.*, 1995, 1561 (*Russ. Chem. Bull.*, 1995, **44**, 1501).
- D. R. Swanson, C. J. Rousset and E. Negishi, *J. Org. Chem.*, 1989, **54**, 3521.
- E. Negishi, D. Choneiri, T. B. Nguyen, D. R. Swanson, N. Suzuki and T. Takahashi, *J. Am. Chem. Soc.*, 1994, **116**, 9751.
- E. Negishi, F. E. Cederbaum and T. Takahashi, *Tetrahedron Lett.*, 1986, **27**, 2829.
- E. Negishi, S. J. Holmes, J. M. Tour, J. A. Miller, F. E. Cederbaum, D. R. Swanson and T. Takahashi, *J. Am. Chem. Soc.*, 1989, **111**, 3336.
- E. Negishi and T. Takahashi, *Acc. Chem. Res.*, 1994, **27**, 124.
- U. M. Dzhemilev, O. S. Vostrikova and A. G. Ibragimov, *Usp. Khim.*, 1986, **55**, 191 (*Russ. Chem. Rev.*, 1986, **55**, 66).
- U. M. Dzhemilev, O. S. Vostrikova and G. A. Tolstikov, *Usp. Khim.*, 1990, **59**, 1972 (*Russ. Chem. Rev.*, 1990, **59**, 1157).
- O. S. Vostrikova, A. G. Ibragimov, R. M. Sultanov and U. M. Dzhemilev, *Metalloorg. Khim.*, 1992, **5**, 782 (*Organomet. Chem. Rev.*, 1992, **5**, 377).

Received: Moscow, 3rd March 1997

Cambridge, 12th June 1997; Com. 7/01771E