

## Atropisomeric and atropdiastereoisomeric 2-substituted 1-aryl-3,5-diphenylpyrroles

Radek Pohl, Jan Pawlas, Stanislav Böhm, Richard Hrabal, Hana Dvořáková and Josef Kuthan\*

<sup>a</sup> Prague Institute of Chemical Technology, 166 28 Prague 6, Czech Republic. Fax: +42 2 311 9919; e-mail: josef.kuthan@vscht.cz

(3,5-Diphenyl-1-*o*-tolyl-1*H*-pyrrol-2-yl)phenylmethanone **3a** and (1-naphthalen-1-yl-3,5-diphenyl-1*H*-pyrrol-2-yl)phenylmethanone **4a** are atropisomeric and their LAH reductions afford mixtures of atropdiastereoisomeric (3,5-diphenyl-1-*o*-tolyl-1*H*-pyrrol-2-yl)phenylmethanols **3b,c** and (1-naphthalen-1-yl-3,5-diphenyl-1*H*-pyrrol-2-yl)phenylmethanols **4b,c**, respectively.

While the atropisomerism caused by restricted rotation around various C–C bonds in biaryls<sup>1</sup> as well as around C–N bonds in heterobiaryls and related heterocycles<sup>2</sup> is well documented by many examples, the axial chirality of 1-arylprrrole derivatives has been observed rarely. Thus, racemic 1-(2-carboxyphenyl)-2,5-dimethyl-1*H*-pyrrole-3-carboxylic acid **1** was resolved<sup>3</sup> in the pioneering studies by Adams and coworkers on possible atropisomeric Knorr pyrroles.<sup>3,4</sup> Many years later, 1-aryl-2,5-dimethylpyrrole-3-carbaldehydes **2** (R = Me, CN, OMe and OCH<sub>2</sub>Ph)<sup>5</sup> have been proven to exhibit axial chirality. We report that (3,5-diphenyl-1-*o*-tolyl-1*H*-pyrrol-2-yl)phenylmethanone **3a** and (1-naphthalen-1-yl-3,5-diphenyl-1*H*-pyrrol-2-yl)phenylmethanone **4a**, which can be easily prepared by extended Decker oxidation<sup>6</sup> of corresponding quaternary pyridinium salts,<sup>†</sup> are also atropisomeric although substituents in their five-membered pyrrole rings may be expected to cause a less hindrance to rotation about the C–N bonds. Indeed, the molecular geometry optimisation for various fixed torsion angles  $\phi$  by the semi-empirical PM3 method<sup>7</sup> predicts the corresponding rotation barriers to be 22.3–27.2 kcal mol<sup>-1</sup> suggesting possible detection of appropriate atropisomers. Hence, <sup>1</sup>H NMR spectra of ketones **3a** and **4a** have been investigated both in the absence and in the presence of the chiral shift reagent Eu(hfc)<sub>3</sub> which caused a typical 1:1 splitting of certain proton signals indicating the racemic character of both of the substrates in C<sub>6</sub>D<sub>6</sub> solutions. For example, two separated 4-H proton and methyl signals clearly indicate this behaviour.<sup>‡</sup>


Similarly, as in a series of axially chiral imidazo[1,2-*a*]pyridinoic ketones,<sup>8</sup> and one 3-chlorobenzoyl derivative of poly-substituted 4-(thien-2-yl)-4*H*-1,2,4-triazoles,<sup>9</sup> the atroposelective reductions of compounds **3a** and **4a** to corresponding secondary alcohols also gave atropdiastereoisomeric products<sup>§</sup> due to a restricted rotation around C(sp<sup>2</sup>)-N(sp<sup>2</sup>) bonds. Thus, LAH with **3a** in diethyl ether at –30 to –25 °C afforded a 7:3 mixture of two (3,5-diphenyl-1-*o*-tolyl-1*H*-pyrrol-2-yl)phenylmethanols **3b,c** while the same procedure with **4a** led analogously to

<sup>†</sup> Details of the preparative procedures will be published elsewhere. IR spectra were taken on 'Nicolet 704' FTIR spectrometer; NMR spectra were recorded on Varian Gemini 300, Bruker AMX 400 and DRX 500 Avance instruments at 298 K.

For **3a**: mp 162–163 °C, yield 83%. <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>)  $\delta$ : 1.69 (s, 3H, *o*-Me), 6.66 (s, 1H, 4-H), 7.00–7.38 (m, 17H, aromatic), 7.60 (d, 2H, *o*-5-Ph). <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>)  $\delta$ : 18.39 (*o*-Me), 111.47 (4-CH), 126.76 (CH), 127.09 (CH), 128.21 (CH), 128.27 (2CH), 128.51 (CH), 128.84 (2CH), 129.04 (2CH), 129.16 (2CH), 129.98 (2CH), 130.21 (CH), 130.65 (CH), 131.16 (CH), 132.35 (CH), 132.55 (CH), 133.65 (C), 135.96 (2C), 136.91 (C), 138.94 (C), 139.17 (C), 140.10 (C), 188.33 (CO). IR (CHCl<sub>3</sub>,  $\nu_{\max}$ /cm<sup>-1</sup>): 1629 (C=O).

For **4a**: mp 156–158 °C, yield 93%. <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>)  $\delta$ : 6.76 (s, H, 4-H), 7.84 (m, 22H, aromatic). <sup>13</sup>C NMR (400 MHz, CDCl<sub>3</sub>)  $\delta$ : 111.73 (4-CH), 123.37 (CH), 125.57 (CH), 126.92 (CH), 127.22 (CH), 127.82 (CH), 128.09 (CH), 128.20 (2CH), 128.28 (CH), 128.59 (2CH), 128.73 (2CH), 128.90 (CH), 129.05 (2CH), 129.49 (CH), 130.06 (2CH), 130.55 (2CH), 131.98 (C), 132.20 (C), 132.49 (CH), 134.46 (C), 135.95 (2C), 136.46 (C), 139.13 (C), 141.47 (C), 188.37 (CO). IR (CHCl<sub>3</sub>,  $\nu_{\max}$ /cm<sup>-1</sup>): 1629 (C=O).

<sup>‡</sup> <sup>1</sup>H NMR  $\delta$  [400 MHz, C<sub>6</sub>D<sub>6</sub>, Eu(hfc)<sub>3</sub> 1:10 molar ratio] for **3a**: 2.05 (s, 3H, *o*-Me, first enantiomer), 2.06 (s, 3H, *o*-Me, second enantiomer), 7.04 (s, 1H, 4-H, first enantiomer), 7.06 (s, 1H, 4-H, second enantiomer); for **4a**: 6.90 (s, 1H, 4-H, first enantiomer), 6.92 (s, 1H, 4-H, second enantiomer).



atropdiastereoisomeric (1-naphthalen-1-yl-3,5-diphenyl-1*H*-pyrrol-2-yl)phenylmethanols **4b,c** in the ratio 8:7.<sup>¶</sup> Pure major atropdiastereoisomers **3b** and **4b** can be separated from the mixtures with their minor counterparts **3c** and **4c** using preparative TLC and/or crystallisation. The atropdiastereoisomers **3b,c** and **4b,c** seem to be quite stable at room and somewhat elevated temperatures both in the solid state and in various solutions. As

<sup>§</sup> All reactions were monitored by HPLC using an Ecom LCP 4000 pump with an UV-VIS detector on Macherey Nagel C-18 guard and common C-18 Nagel columns, eluent MeOH–H<sub>2</sub>O 4:1. Preparative TLC was performed on 20×20 cm plates of Aldrich TLC high purity grade silica gel (15 g) with a gypsum binder and a fluorescent indicator; eluent PE–DCM 5:1. All products gave satisfactory elemental analyses.

<sup>¶</sup> Atropdiastereoisomeric mixture of **3b** and **3c**: HPLC (MeOH–H<sub>2</sub>O 7:3, two peaks 111.6 and 121.5 min), mp 129–130 °C, yield 89%. <sup>1</sup>H NMR (300 MHz, [<sup>2</sup>H<sub>6</sub>]DMSO)  $\delta$ : 1.14 (s, 3H, *o*-Me), 1.82 (s, 3H, *o*-Me), 5.70 (s, 1H, H-6 ?), 5.93 (d, OH-6 and H-6 ?), 6.59 (s, 1H, 4-H), 6.70–7.78 (m, aromatic). IR (KBr,  $\nu_{\max}$ /cm<sup>-1</sup>): 3526 (OH).

Major atropdiastereoisomer **3b**: HPLC (MeOH–H<sub>2</sub>O 7:3, one peak 111.6 min); mp 133–137 °C (diethyl ether–heptane). <sup>1</sup>H NMR (300 MHz, [<sup>2</sup>H<sub>6</sub>]DMSO)  $\delta$ : 1.14 (s, 3H, Me-7'), 5.93 (m, 2H, OH-6 and H-6), 6.57 (s, 1H, H-4), 6.70–7.76 (m, 19H, aromatic). <sup>13</sup>C NMR (75.4 MHz, [<sup>2</sup>H<sub>6</sub>]DMSO)  $\delta$ : 16.88 (Me, C-7'), 65.91 (CH, C-6), 109.07 (CH, C-4), 124.53 (C), 125.10 (2CH), 125.62 (2CH), 125.95 (CH), 126.03 (CH), 126.40 (CH), 127.03 (2CH), 127.20 (2CH), 128.17 (2CH), 128.42 (2CH), 128.80 (2CH), 129.90 (CH), 131.31 (CH), 132.90 (C), 133.35 (C), 134.43 (C), 136.37 (2C), 137.59 (C), 142.85 (C).

Atropdiastereoisomeric mixture of **4b** and **4c**: HPLC (MeOH–H<sub>2</sub>O 4:1 or 7:3, a single broad peak 26.3 or 160.9 min, respectively), mp 165–171 °C, yield 94%. <sup>1</sup>H NMR (300 MHz, [<sup>2</sup>H<sub>6</sub>]DMSO) 5.73 (d, 1H, OH-6), 5.86 (dd, 2H, H-6 and H'-6), 5.93 (d, 1H, OH-6), 6.69 (s, 1H, H-4), 6.74 (s, 1H, H-4), 6.44–7.96 (m, 22H, aromatic). IR ( $\nu_{\max}$ /cm<sup>-1</sup>): 3545 (OH).

Major atropdiastereoisomer **4b**: HPLC (MeOH–H<sub>2</sub>O 7:3, one peak 160.2 min), mp 174–178 °C. <sup>1</sup>H NMR (300 MHz, [<sup>2</sup>H<sub>6</sub>]DMSO)  $\delta$ : 5.86 (d, 1H, H-6, *J* 3.8 Hz), 5.93 (d, 1H, OH-6, *J* 3.8 Hz), 6.69 (s, 1H, H-4), 6.44–7.96 (m, 22H, aromatic). <sup>13</sup>C NMR (75.4 MHz, [<sup>2</sup>H<sub>6</sub>]DMSO)  $\delta$ : 65.88 (CH, C-6), 109.39 (CH, C-4), 123.18 (CH), 124.70 (CH), 124.77 (CH), 124.96 (CH), 125.75 (CH), 126.12 (CH), 126.19 (CH), 126.44 (CH), 126.54 (CH), 127.17 (CH), 127.44 (CH), 128.03 (CH), 128.44 (CH), 128.86 (CH), 129.05 (CH), 130.78 (C), 132.97 (C), 133.13 (C), 134.92 (C), 135.23 (C), 135.63 (C), 136.32 (C), 142.17 (C). IR ( $\nu_{\max}$ /cm<sup>-1</sup>): 3545.

expected, the presence of the methyl group in the molecules of **3a–c** should be a necessary condition for the axial chirality behaviour. In fact, the formerly reported<sup>10</sup> 1-phenyl derivative **3d** is not atropisomeric under the same conditions. Quantum-chemical calculations and experiments with other **3a**-like ketones as well as an extended study on the stereochemistry of **3b**- and **3c**-like alcohols are in progress.

This work was supported by the Grant Agency of the Czech Republic (grant no. 203/96/0497) and the Czech Ministry of Education (grant no. 005/G).

## References

- (a) M. Oki, in *Topics in Stereochemistry*, eds. N. L. Allinger, E. L. Eliel and S. H. Wilen, Wiley, New York, 1983, vol. 14, p. 1; (b) E. L. Eliel, *Stereochemistry of Organic Compounds*, Wiley, New York, 1994.
- (a) Z. Aksač, E. Pinar and S. Içli, *Org. Magn. Reson.*, 1983, **21**, 548; (b) A. Mannschreck, H. Koller, G. Stühler, M. A. Davies and J. Traber, *Eur. J. Med. Chem. Chim. Ther.*, 1984, **19**, 381; (c) C. Roussel, M. Adjimi, A. Chemlal and A. Djafri, *J. Org. Chem.*, 1988, **53**, 5076; (d) M. Mintas, V. Mihaljević, H. Koller, D. Schuster and A. Mannschreck, *J. Chem. Soc., Perkin Trans. 2*, 1990, 619; (e) I. Dogan, N. Pustet and A. Mannschreck, *J. Chem. Soc., Perkin Trans. 2*, 1993, 1557; (f) K. Saito, M. Yamamoto and K. Yamada, *Tetrahedron*, 1993, **49**, 4549; (g) D. P. Curran, H. Qi, S. J. Geib and N. C. DeMello, *J. Am. Chem. Soc.*, 1994, **116**, 3131; (h) C. Roussel and C. Suteu, *J. Chromatogr. A*, 1997, **761**, 129; (i) X. Dai, A. Wong and S. C. Virgil, *J. Org. Chem.*, 1998, **63**, 2597; (j) O. Kitagawa, H. Izawa, K. Sato, A. Dobashi, T. Taguchi and M. Shiro, *J. Org. Chem.*, 1998, **63**, 2634.
- L. H. Bock and R. Adams, *J. Am. Chem. Soc.*, 1931, **53**, 374.
- (a) C. Chang and R. Adams, *J. Am. Chem. Soc.*, 1931, **53**, 2353; (b) L. H. Bock and R. Adams, *J. Am. Chem. Soc.*, 1931, **53**, 3519.
- (a) J. Vorkapić-Furač, M. Mintas, T. Burgemeister and A. Mannschreck, *J. Chem. Soc., Perkin Trans. 2*, 1989, 713; (b) J. Vorkapić-Furač, M. Mintas, F. Kastner and A. Mannschreck, *J. Heterocycl. Chem.*, 1992, **29**, 327.
- J. Kuthan, *Heterocycles*, 1994, **37**, 1347.
- J. J. P. Stewart, *Comput. Chem.*, 1989, **10**, 209, 221.
- (a) R. Kubík, J. Němeček, S. Böhm, M. Hradilek and J. Kuthan, *Mendeleev Commun.*, 1995, 29; (b) R. Kubík, S. Böhm and J. Kuthan, *Collect. Czech. Chem. Commun.*, 1996, **61**, 1018.
- N. Marubayashi, T. Ogawa, M. Moriwaki and M. Haratake, *Chem. Pharm. Bull.*, 1992, **40**, 220.
- P. Nesvadba and J. Kuthan, *Collect. Czech. Chem. Commun.*, 1982, **47**, 1494.

Received: Cambridge, 28th September 1998

Moscow, 10th November 1998; Com. 8/07516F