

# Thermodynamic dissociation constants for the $[\text{Cu}(\text{py})_4(\text{NO}_3)_2] \cdot 2\text{py}$ clathrate and $[\text{Cu}(\text{py})_4(\text{NO}_3)_2]$ complex


Elissa A. Ukraintseva, Dmitriy V. Soldatov, Yuri A. Dyadin,\* Vladimir A. Logvinenko and Eugeny V. Grachev

Institute of Inorganic Chemistry, Siberian Branch of the Russian Academy of Sciences, 630090 Novosibirsk, Russian Federation.  
Fax: +7 3832 34 4489; e-mail: clat@che.nsk.su

A comparison of the determined thermodynamic dissociation parameters for  $[\text{Cu}(\text{py})_4(\text{NO}_3)_2] \cdot 2\text{py}$  with those for analogous Mn, Co, Ni and Zn clathrates indicates the stability of the host molecules as the main contribution to the stability of the resulting clathrates.

Supramolecular compounds  $[\text{M}(\text{py})_4(\text{NO}_3)_2] \cdot 2\text{py}$  ( $\text{M} = \text{Mg}, \text{Mn}, \text{Co}, \text{Ni}, \text{Cu}, \text{Zn}$  or  $\text{Cd}$ ) form in the pyridine (py)–metal(II) nitrate systems.<sup>1–7</sup> There are two types of pyridine molecules in the compounds: ligated to the metal atom (ligand pyridine) and located inside crystal framework cavities (guest pyridine). Our studies of the Cu system revealed a surprisingly high stability of the  $[\text{Cu}(\text{py})_4(\text{NO}_3)_2] \cdot 2\text{py}$  clathrate as well as the  $[\text{Cu}(\text{py})_4(\text{NO}_3)_2]$  host complex formed in the system (Table 1).<sup>†</sup> The clathrate endures heating of up to 145 °C melting incongruently to produce the solid host complex at this temperature. The decomposition of its structural analogues formed in the Mn, Co, Ni, Zn and Cd systems occurs at lower temperatures without the stage of formation of the host complexes which undergo further dissociation. The resulting tris(pyridine) complexes,  $[\text{M}(\text{py})_3(\text{NO}_3)_2]$ , are isomorphous to one another<sup>2,12,13</sup> as well as to  $[\text{Cu}(\text{py})_3(\text{NO}_3)_2]$ <sup>12</sup> which forms during the  $[\text{Cu}(\text{py})_4(\text{NO}_3)_2]$  decomposition. From these data, we assumed that it is the strength of host complex coordination bonds that mainly contributes to the overall stability of the resulting clathrate phases, rather than the host-to-guest complementarity, which is a decisive factor for classical clathrates.<sup>14</sup> To provide support for this assumption, it was necessary to compare thermodynamic dissociation parameters for this clathrate series. As such data had been available for all the clathrates of transition metal complex hosts except the Cu system,<sup>9,11,15,16</sup> we determined the pyridine vapour pressure over the title compounds as a function of temperature.

Synthesis and analysis of the compounds were described earlier.<sup>17</sup> The stoichiometry of dissociation was studied with the quasi-equilibrium thermogravimetry method.<sup>‡</sup> The numerical



**Figure 1** Mass loss curves for  $[\text{Cu}(\text{py})_4(\text{NO}_3)_2] \cdot 2\text{py}$ . Sample, 150 mg; atmosphere, air; heating, quasi-isobaric-quasi-isothermal;  $q = 0.3 \text{ mg min}^{-1}$ ; sample holders: (1) plate, (2) open, (3) closed and (4) conic crucibles.


<sup>†</sup> Melting points given in text and Table 1 were measured by DTA using samples sealed in glass ampoules. Heating rate:  $1 \text{ K min}^{-1}$ . Accuracy of phase transition temperatures 0.9 K or better. Note that under other conditions the compounds decompose at lower temperatures with the loss of pyridine (Figure 1).

**Table 1** Melting points (°C) of some compounds in the pyridine-metal(II) nitrate systems.


| M  | $[\text{M}(\text{py})_3(\text{NO}_3)_2]$ | $[\text{M}(\text{py})_4(\text{NO}_3)_2]$ | $[\text{M}(\text{py})_4(\text{NO}_3)_2] \cdot 2\text{py}$ | Reference |
|----|--|--|---|-----------|
| Mn | 159                                      | unknown                                  | 81 (decomp.)  | 2         |
| Co | 160                                      | unknown                                  | 88 (decomp.)  | 8 and 9   |
| Ni | 165                                      | unknown                                  | 108   | 8 and 9   |
| Cu | 172                                      | 175                                      | 145 (decomp.)   | this work |
| Zn | 131                                      | unstable                                 | 62 (decomp.)  | 10        |
| Cd | 165                                      | unstable                                 | 106 (decomp.)   | 11        |

data on the equilibrium pyridine vapour pressure over the title compounds as a function of temperature were measured using a strain technique.<sup>§</sup>

The mass loss curves for  $[\text{Cu}(\text{py})_4(\text{NO}_3)_2] \cdot 2\text{py}$  are shown in Figure 1. The clathrate dissociates with formation of the host complex:



The complex dissociates further:



The data on the vapour pressure are shown in Figure 2. The temperature dependence of the pyridine vapour pressure during the clathrate dissociation exhibits a bend at approximately 333 K.<sup>¶</sup> The numerical data on the pyridine vapour pressure over the title compounds are approximated by the equations  $\log P = A - B/T$ . The coefficients and the calculated thermodynamic dissociation parameters are summarised in Table 2.

The parameters for the total transformation of the clathrate into the tris(pyridine) complex calculated from these data are given in Table 3 along with corresponding parameters for other

<sup>‡</sup> MOM Q-1500-D derivatograph (Hungary). The mass loss curves were recorded at a rate of  $0.3 \text{ mg min}^{-1}$ , the mass of the samples was 150 mg. The set of standard holders provided the following pressures of the product gas: 0.01 (a plate crucible), 0.05 (an open crucible), 0.2 (a closed crucible) and 0.8 atm (a conic crucible).

<sup>§</sup> The vapour pressure was measured by a static method using a Pyrex spoon-type membrane as a null manometer.<sup>19</sup> The membrane sensitivity varied from 0.05 to 0.1 Torr in different experimental series. The accuracy of thermostating and temperature measurements was 0.05 K. The volume of the reaction vessel was ca. 50 ml.


The membrane chamber was filled with substances in a dry box and sealed after evacuating. For  $[\text{Cu}(\text{py})_4(\text{NO}_3)_2] \cdot 2\text{py}$ , two experimental runs over the temperature range 304–378 K were performed. Clathrate portions of 143 and 719 mg were used in the first and second runs, respectively, with the same reaction vessel volume. No significant difference between the data obtained in these two runs was observed. The clathrate dissociation is completely reversible; the equilibrium pressure established well both in the forward and backward directions. For  $[\text{Cu}(\text{py})_4(\text{NO}_3)_2]$ , one run over the range 337–392 K was performed (the pressure was < 1 Torr at lower temperatures). The complex dissociation is also reversible although the backward reaction is significantly slower.

<sup>¶</sup> The bend corresponds to a polymorphous transformation of the clathrate. Possibly, it is analogous to the transformation of  $[\text{Cd}(\text{py})_4(\text{NO}_3)_2] \cdot 2\text{py}$  at 221 K.<sup>7</sup>

**Table 2** Coefficients of the  $\log P = A - B/T$  equations and calculated thermodynamic parameters of reactions (1) and (2).

| Reaction | Number of experimental points | T range/K | A        | B/1000  | $H_{av}^{\circ}/\text{kJ mol}^{-1}$ | $S_{av}^{\circ}/\text{J mol}^{-1} \text{ K}^{-1}$ | $G_{298}^{\circ}/\text{kJ mol}^{-1}$ |
|----------|-------------------------------|-----------|----------|---------|-------------------------------------|---|--------------------------------------|
| (1)      | 15                            | 304–3 33  | 13.2(2)  | 3.90(5) | 149(2)                              | 394(6)  | 31.8(9)                              |
| (1)      | 17                            | 333–3 78  | 10.43(4) | 2.99(2) | 114.4(6)                            | 289(2)  | 28.2(8)                              |
| (2)      | 11                            | 338–3 92  | 11.3(1)  | 3.66(4) | 70.1(7)                             | 162(2)  | 21.9(9)                              |

clathrates of the  $[\text{M}(\text{py})_4(\text{NO}_3)_2] \cdot 2\text{py}$  series. Due to the isomorphism of both clathrates and tris(pyridine) complexes, the differences in packing energies are insignificant, and it is the strength of the metal-pyridine bond that should have a predominant influence on differences in listed thermodynamic constants for the following process:



For the Cu compound, reaction (3) is a sum of (1) and (2), while in other systems the intermediate host complex  $[\text{M}(\text{py})_4(\text{NO}_3)_2]$  does not form an individual phase.

The dependence of the clathrate dissociation parameters on the nature of metal in the host complex (Table 3) confirms this hypothesis. The thermodynamic stability of the clathrates in the series does not correlate with packing coefficients<sup>†</sup>,<sup>20</sup> or other structural parameters; a correlation was usually found for clathrate compounds.<sup>14</sup> At the same time, the order of stability of these clathrate phases follows the general stability sequence for complexes (Irwing-Williams sequence<sup>21</sup>)  $\text{Mn} < \text{Fe} < \text{Co} < \text{Ni} < \text{Cu} > \text{Zn}$ .

This demonstrates how the strengthening of the coordination bonds of the host complex results in a progressive rise in the clathrate phase stability. Note that in the majority of systems the host complex is insufficiently stable to form its own, non-clathrate, phase. This phenomenon, when the molecules unable to create a phase can be stabilised by non-covalent contact with molecules of another type, was referred to as contact stabilisation of molecules.<sup>22</sup> So far we have determined the measure of instability of a molecule as the inability to have its own phase. Here, we quantitatively determine this measure of instability (Table 3). Only in the Cu system, the host molecule can form a stable host phase. This qualitative change manifests itself as a steep quantitative rise in the thermal and thermodynamic stability of the clathrate formed by the complex.


The conventional approach considers the stability of clathrates as a result of the complementarity between host and guest molecules. Indeed, this approach is quite justified provided the participating molecules are stable enough to prevent dissociation during all processes under consideration. It is evident, however, that the overall stability of clathrates depends not only on inter-

**Table 3** The comparison of the pyridine-metal(II) nitrate systems. Packing coefficients ( $k_{\text{pack}}$ ) of the isomorphous clathrates  $[\text{M}(\text{py})_4(\text{NO}_3)_2] \cdot 2\text{py}$  and thermodynamic parameters of reaction (3).

| M  | $k_{\text{pack}}$ | T range/K                                      | $H_{av}^{\circ}/\text{kJ mol}^{-1}$ | $S_{av}^{\circ}/\text{J mol}^{-1} \text{ K}^{-1}$ | $G_{298}^{\circ}/\text{kJ mol}^{-1}$ | Reference              |
|----|-------------------|--|-------------------------------------|---|--------------------------------------|------------------------|
| Mn | 0.660(3)          | 292–35 2                                       | 160.8(9)                            | 424(3)  | 34(2)                                | 15                     |
| Co |                   | 293–36 0                                       | 178.4(5)                            | 470(2)  | 38.3(7)                              | 9                      |
| Ni | 0.672(3)          | 293–34 5                                       | 194(1)                              | 505(4)  | 43(2)                                | 9                      |
| Cu | 0.669(3)          | 304–33 3 <sup>a</sup><br>333–37 8 <sup>a</sup> | 219(3)<br>185(1)                    | 556(8)<br>451(4)                                  | 54(2)<br>50(2)                       | this work<br>this work |
| Zn | 0.663(3)          | 300–33 5                                       | 175(2)                              | 475(6)  | 33(3)                                | 16                     |
| Cd | 0.659(3)          | 290–36 0                                       | 165(1)                              | 426(3)  | 38(1)                                | 11                     |

<sup>a</sup>For the Cu system, reaction (3) is the sum of (1) and (2), which were studied in different temperature ranges (Table 2); here, we assume that the dissociation parameters for the  $[\text{Cu}(\text{py})_4(\text{NO}_3)_2]$  complex in the indicated temperature interval are the same as in the temperature range studied.

<sup>†</sup>The volumes of the host and pyridine molecules were calculated using X-ray structural data for the clathrates ( $\text{Mn}$ ,<sup>2</sup>  $\text{Ni}$ ,<sup>4</sup>  $\text{Cu}$ ,<sup>5</sup>  $\text{Zn}$ <sup>6</sup> and  $\text{Cd}$ <sup>7</sup>). As the structures of the Cu and Zn clathrates were determined at low temperatures, the unit cell parameters measured at room temperature were taken from ref. 3.

**Figure 2** Temperature dependence of the pyridine vapour pressure over (1)  $[\text{Cu}(\text{py})_4(\text{NO}_3)_2] \cdot 2\text{py}$  and (2)  $[\text{Cu}(\text{py})_3(\text{NO}_3)_2]$  (experimental data) and (3) over liquid pyridine<sup>16</sup> on the  $\log P - 1000/T$  coordinates ( $P/\text{Torr}$ ,  $T/\text{K}$ ).

molecular interactions but also on the stability of the component molecules. Therefore, the latter factor may be decisive, and this is the case with the clathrate series under consideration.

This work was supported by the Federal Special Programme 'State Foundation for the Integration of Higher Education and Fundamental Science in 1997–2000' (grant no. 274).

## References

- D. V. Soldatov and J. A. Ripmeester, *Supramol. Chem.*, 1998, **9**, 175 and references therein.
- D. V. Soldatov and J. Lipkowski, *Zh. Strukt. Khim.*, 1998, **39**, 293 [*J. Struct. Chem. (Engl. Transl.)*, 1998, **39**, 238].
- D. V. Soldatov and J. Lipkowski, *Zh. Strukt. Khim.*, 1995, **36**, 1070 [*J. Struct. Chem. (Engl. Transl.)*, 1995, **36**, 979].
- D. V. Soldatov and J. Lipkowski, *Zh. Strukt. Khim.*, 1995, **36**, 912 [*J. Struct. Chem. (Engl. Transl.)*, 1995, **36**, 830].
- G. Beurskens, C. F. Martens, R. J. M. Nolte, P. T. Beurskens and J. M. M. Smits, *J. Chem. Crystallogr.*, 1995, **25**, 425.
- P. Losier and M. J. Zaworotko, *J. Chem. Crystallogr.*, 1996, **26**, 277.
- D. V. Soldatov, B. A. Kolesov, J. Lipkowski and Yu. A. Dyadin, *Zh. Strukt. Khim.*, 1997, **38**, 976 [*J. Struct. Chem. (Engl. Transl.)*, 1997, **38**, 819].
- P. Tomasik and Z. Ratajczak, *Pyridine-Metal Complexes*, in *The Chemistry of Heterocyclic Compounds*, eds G. R. Newkome and L. Strekowski, Wiley, New York, 1985, vol. 14, pt. 6, pp. 1118, 1452.
- E. A. Ukraintseva, D. V. Soldatov, V. A. Logvinenko and Yu. A. Dyadin, *Mendeleev Commun.*, 1997, 102.
- D. V. Soldatov, V. A. Logvinenko and Yu. A. Dyadin, *Zh. Neorg. Khim.*, 1995, **40**, 324 (*Russ. J. Inorg. Chem.*, 1995, **40**, 309).
- D. V. Soldatov, Yu. A. Dyadin, E. A. Ukraintseva, B. A. Kolesov and V. A. Logvinenko, *J. Inclusion Phenom.*, 1996, **26**, 269.

- 12 A. F. Cameron, D. W. Taylor and R. H. Nuttall, *J. Chem. Soc., Dalton Trans.*, 1972, 1603.
- 13 A. F. Cameron, D. W. Taylor and R. H. Nuttall, *J. Chem. Soc., Dalton Trans.*, 1972, 1608.
- 14 *Comprehensive Supramolecular Chemistry*, eds. J. L. Atwood, J. E. D. Davies, D. D. MacNicol and F. Vögtle, Pergamon, Oxford, 1996.
- 15 E. A. Ukraintseva, D. V. Soldatov and Yu. A. Dyadin, *Collected Abstracts of the Xth International Symposium on Molecular Recognition and Inclusion*, Warsaw, 1998, p. P-4.
- 16 E. A. Ukraintseva, D. V. Soldatov and Yu. A. Dyadin, *Zh. Neorg. Khim.*, 1997, **42**, 283 (*Russ. J. Inorg. Chem.*, 1997, **42**, 229).
- 17 D. V. Soldatov, Yu. A. Dyadin, J. Lipkowski and A. G. Ogienko, *Mendeleev Commun.*, 1997, 11.
- 18 F. Paulik and J. Paulik, *J. Therm. Anal.*, 1973, **5**, 253.
- 19 E. A. Ukraintseva, Yu. A. Dyadin, N. V. Kislykh, V. A. Logvinenko and D. V. Soldatov, *J. Inclusion Phenom.*, 1995, **23**, 23.
- 20 A. I. Kitaigorodsky, *Molecular Crystals and Molecules*, Academic Press, New York, 1973, p. 18.
- 21 H. Irving and R. J. P. Williams, *J. Chem. Soc.*, 1953, 3192.
- 22 (a) Yu. A. Dyadin and N. V. Kislykh, *Mendeleev Commun.*, 1991, 134;  
(b) Yu. A. Dyadin, D. V. Soldatov, V. A. Logvinenko and J. Lipkowski, *J. Coord. Chem.*, 1996, **37**, 63.

Received: 27th November 1998; Com. 98/1406 (8/09471C)