

Simple synthesis of natural polyketides, 2-dodecanoyl-5-hydroxycyclohexane-1,3-dione and 2-dodecanoylresorcinol

Vladimir G. Zaitsev,* Pavel M. Philipchenko and Fedor A. Lakhvich

Institute of Bioorganic Chemistry, National Academy of Sciences of Belarus, 220141 Minsk, Belarus.
Fax: +375 172 63 7274; e-mail: prostan@ns.iboch.ac.by

The title compounds were synthesised in two and three steps, respectively, starting from 5-hydroxycyclohexane-1,3-dione.

2-Dodecanoyl-5-hydroxycyclohexane-1,3-dione **1** was identified¹ as the major component of the secretions of *Stephanitis takeyai*. It was prepared by a complex multistage synthesis including the construction of the cyclohexane ring containing a phenyldimethylsilyl substituent as a masked hydroxy group.² The authors reported the easy aromatization of hydroxytriketone **1** due to the known susceptibility of α -hydroxyketones to dehydration.

Recently, we reported the first synthesis of 5-hydroxycyclohexane-1,3-dione **2**.³ We have found that 5-hydroxydiketone **2** exists in solution mainly in the enolic form, and it takes part in reactions through its enol group. It gives little or no by-product aromatics in the absence of an acid. Based on this finding, we carried out a simple two-step synthesis of target compound **1** by the method developed earlier⁴ for regioisomeric 2-acyl-4-hydroxycyclohexane-1,3-diones. Acylation of 5-hydroxydiketone **2** with dodecanoyl chloride proceeds regioselectively, affording enolacrylate **3**. The latter was then subjected to a smooth O–C isomerization into triketone **1**[†] under the CN⁻ catalysis.


This method can be used as a general procedure for the synthesis of other 2-acyl-5-hydroxycyclohexane-1,3-diones. Moreover, these 5-hydroxyketones are the immediate synthetic precursors of natural bioactive 2-acylresorcinols.⁵ Thus, hydroxytriketone **1** was entirely converted into 2-dodecanoylresorcinol **4**[‡] when treated with a mineral acid.

This work was supported by the Byelarusian Foundation for Basic Research (grant no. X97-109).

[†] 2-Dodecanoyl-5-hydroxycyclohexane-1,3-dione **1**. To a stirred solution of 0.39 g (3 mmol) of anhydrous 5-hydroxycyclohexane-1,3-dione **2** (mp 95–96 °C) and 0.25 ml (3.1 mmol) of pyridine in 30 ml of dioxane, 0.60 ml (2.5 mmol) of dodecanoyl chloride in 20 ml of dioxane was added dropwise, over 1 h at room temperature. The solvent was evaporated *in vacuo*, and 20 ml of CHCl₃ was added to the residue. The solution was washed with 0.2 M HCl, H₂O, brine, and then dried over MgSO₄. The crude product after solvent evaporation was recrystallised from hexane giving 0.71 g (92%) of 3-dodecanoyl-5-hydroxycyclohex-2-en-1-one **3**. Mp 43–44 °C. ¹H NMR (200 MHz, CDCl₃) δ : 0.88 (t, 3H, Me, *J* 6.5 Hz), 1.27 (m, 16H, 4'-1'-CH₂), 1.67 (m, 2H, 3'-CH₂), 2.48 (t, 2H, 2'-CH₂, *J* 7.5 Hz), 2.55 and 2.64 (2dd, 2H, 4,6-H_a, *J* 6.5 and 17 Hz), 2.70 (dd, 1H, 4- or 6-H_c, *J* 4 and 17 Hz), 2.88 (dd, 1H, 4- or 6-H_c, *J* 4 and 17 Hz), 4.43 (m, 1H, 5-H), 5.98 (s, 1H, 2-H). IR (KBr, ν /cm⁻¹): 3440 (br.), 1765, 1680, 1660, 1480, 1420, 1380, 1145, 1135, 1115, 1070.

To a solution of 0.17 g (0.55 mmol) of enolacrylate **3** and 0.24 ml (2.5 mmol) of Et₃N in 30 ml of acetonitrile, 0.06 ml (0.6 mmol) of acetone cyanohydrin was added. The reaction mixture was allowed to stand for 3 h at room temperature, then the solvent was evaporated *in vacuo*, and the residue was worked up as above giving, after recrystallization from hexane, 0.16 g (94%) of 2-dodecanoyl-5-hydroxycyclohexane-1,3-dione **1**. Mp 55–56 °C. Physical and chemical characteristics of synthetic **1** are the same as those published.²

[‡] 2-Dodecanoylresorcinol **4** was obtained by dehydration in an acid medium (2 drops of conc. HCl in 20 ml of acetone) of crude hydroxytriketone **1** synthesised as described above from 0.30 g (0.97 mmol) of enolacrylate **3**. After completing the dehydration reaction (3 h, control by TLC), the solvent was evaporated. The residue was dissolved in CHCl₃, washed with H₂O and brine, dried over MgSO₄, and the solvent was evaporated giving, after recrystallization of the residue from hexane, 0.25 g (88%) of 2-dodecanoylresorcinol **4**. Mp 85–86 °C. Physical and chemical characteristics of synthetic **4** are the same as those published.⁶



References

- 1 J. E. Oliver, W. R. Lusby and J. W. Neal, Jr., *J. Chem. Ecol.*, 1990, **16**, 2243.
- 2 J. E. Oliver, R. M. Waters and W. R. Lusby, *Tetrahedron*, 1990, **46**, 1125.
- 3 V. G. Zaitsev and F. A. Lakhvich, *Mendeleev Commun.*, 1998, 20.
- 4 V. G. Zaitsev, G. I. Polozov and F. A. Lakhvich, *Tetrahedron*, 1994, **50**, 6377.
- 5 (a) R. A. Jurenka, J. W. Neal, Jr., R. W. Howard, J. E. Oliver and G. I. Blomquist, *Comp. Biochem. Physiol.*, 1989, **93C**, 253; (b) Y. Tsuda, Sh. Hosoi and Y. Goto, *Chem. Pharm. Bull.*, 1991, **39**, 18; (c) V. G. Zaitsev and F. A. Lakhvich, *Mendeleev Commun.*, 1995, 224.
- 6 K. K. Purushotaman, A. Sarada and J. D. Connolly, *J. Chem. Soc., Perkin Trans. 1*, 1977, 587.

Received: 18th December 1999; Com. 98/1414