

Synthesis of nitro-substituted benzoannelated seven-membered heterocycles from trinitrotoluene

Natalya B. Chernysheva, Alexander V. Samet, Viktor N. Marshalkin, Valery A. Polukeev and Victor V. Semenov*

N. D. Zelinsky Institute of Organic Chemistry, Russian Academy of Sciences, 119991 Moscow, Russian Federation.


Fax: +7 095 135 5328

10.1070/MC2001v011n03ABEH001426


Dibenzo[*b,f*]oxepine, dibenzo[*b,f*]-1,4-oxazepinone and benzo[*f*]naphtho[2,1-*b*]-1,4-oxazepine were synthesised starting from 2,4,6-trinitrotoluene.

While investigating the utilization of 2,4,6-trinitrotoluene (TNT), it was found to be a promising starting compound for the synthesis of benzoannelated five-membered heterocycles (indoles, indazoles, benzothiophenes and benzo[*d*]isothiazoles).^{1–4}

This communication deals with the synthesis of seven-membered N- and O-containing benzoannelated heterocycles with nitro groups at the benzene ring from TNT. These compounds are of special interest because of their pharmaceutical activity.^{5,6} Thus, the antidepressant Sintamil probably possesses a structure of dibenzo[*b,f*]-1,4-oxazepinone:⁷



Reactions of TNT with salicylaldehyde and 1-nitroso-2-naphthol afforded dibenzo[*b,f*]oxepine **1** and benzo[*f*]naphtho[2,1-*b*]-1,4-oxazepine **2**, respectively (Scheme 1).[†]



Scheme 1 Reagents and conditions: i, morpholine, reflux, 3 h, 75%; ii, Et₃N, MeCN, 5 min, 31%.


[†] ¹H NMR spectra were recorded in [2H₆]DMSO on a Bruker DRX500 spectrometer (500.13 MHz), mass spectra were measured on a Kratos MS-30 instrument. ¹H NMR spectra for the reaction products of TNT with salicylaldehydes, 1-nitroso-2- and 2-nitroso-1-naphthols and the ¹H NMR spectra of the products of nucleophilic substitution in oxazepines **2** and **4** are placed on the web-site <http://www.chemical-block.com>.

1,3-Dinitrodibenzo[*b,f*]oxepine 1: to a mixture of 10 mmol (2.27 g) of TNT and 35 mmol (4.27 g) salicylaldehyde, morpholine (10 mol% on an aldehyde basis) was added, and the mixture was heated for 3 h at 135 °C. Then, the reaction mixture was cooled, dissolved in hot AcOH and filtered. The filtrate was cooled; the precipitate was filtered off and dried. Yield 2.13 g (75%), yellowish brown crystals, mp 148–150 °C. ¹H NMR, δ: 7.02 (d, 1H, *J* 12.1 Hz), 7.29 (m, 2H), 7.43 (d, 1H, *J* 8.1 Hz), 7.50 (m, 2H), 8.51 (s, 1H, H-2 or H-4), 8.59 (s, 1H, H-2 or H-4). MS, *m/z* (%): 284 (M⁺, 15), 255 (59), 225 (53), 179 (100), 163 (61).

1,3-Dinitrobenzo[*f*]naphtho[2,1-*b*]-1,4-oxazepine 2: to a solution of 0.346 g (2 mmol) of 1-nitroso-2-naphthol and 0.454 g (2 mmol) of TNT in 3 ml of MeCN, 0.20 g (2 mmol) of Et₃N was added with stirring. After 5 min, the precipitate was filtered off, washed with MeOH and acetone and dried. Yield 0.208 g (31%), deeply orange crystals, mp 276–278 °C. ¹H NMR, δ: 7.55–7.65 (m, 3H), 7.92 (m, 2H), 8.32 (d, 1H, *J* 8.4 Hz), 8.60 (s, 1H, H-2 or H-4), 8.68 (s, 1H, H-2 or H-4), 9.19 (s, 1H, H-13). MS, *m/z* (%): 335 (M⁺, 100), 243 (68), 231 (43), 214 (41).

Probably, in both cases the methyl group of TNT reacts first (with the C=O or N=O group of the reagent molecule) followed by cyclization of the intermediate product *via* intramolecular nucleophilic replacement of the *o*-nitro group by the OH group. Substituted salicylaldehydes, as well as 2-nitroso-1-naphthol, react in a similar way.

Dibenzo[*b,f*]-1,4-oxazepine **4** was obtained from 2,4,6-trinitrobenzoyl chloride (readily prepared from TNT).[‡] In this case, intermediate amide **3** can be isolated, and its cyclization to oxazepinone **4** occurs slowly (Scheme 2).



Scheme 2 Reagents and conditions: i, C₆H₆, reflux, 8 h, 81%; ii, aqueous NH₃, EtOH–MeCN, 50 °C, 60 h, 91%.

Dinitro-substituted compounds **1**, **2** and **4** react with O- and S-nucleophiles to yield nucleophilic substitution products in which one or two nitro groups are replaced. These results will be published elsewhere.

References

- A. V. Samet, V. N. Marshalkin, S. G. Zlotin, V. V. Semenov, A. A. Gakh and A. C. Buchanan III, *218th ACS National Meeting*, New Orleans, 1999, p. 47.
- S. G. Zlotin, P. G. Kisilitsin, A. V. Samet, E. A. Serebryakov, L. D. Konyushkin, V. V. Semenov, A. C. Buchanan III and A. A. Gakh, *J. Org. Chem.*, 2000, **65**, 8430.

[‡] **2,4,6-Trinitrobenzoic acid (2-hydroxyphenyl)amide 3:** prepared in 81% yield according to the general procedure for 2,4,6-trinitrobenzamides.² Grey powder, mp 201–203 °C. ¹H NMR, δ: 6.80–7.00 (m, 3H), 8.05 (d, 1H, *J* 8.3 Hz), 9.10 (s, 2H), 9.70 (br. s, 1H, OH), 10.35 (br. s, 1H, NH). MS, *m/z* (%): 348 (M⁺, 5), 153 (69), 126 (81), 108 (99), 80 (83), 79 (100).

1,3-Dinitrodibenzo[*b,f*]-1,4-oxazepin-11(10H)-one 4: to a solution of 0.348 g (1 mmol) of amide **3** in a mixture of EtOH (2 ml) and MeCN (2 ml), 7 drops of 25% aqueous NH₃ were added, and the mixture was heated for 60 h at 50 °C until the starting amide disappeared (TLC monitoring). Then, the precipitate was filtered off, washed with EtOH and dried. Yield 0.275 g (91%), black needles (orange needles after chromatography on silica gel; eluent, benzene–EtOAc), mp 289–290 °C. ¹H NMR, δ: 7.15–7.30 (m, 3H), 7.48 (d, 1H, H-9, *J* 8.5 Hz), 8.49 (s, 1H, H-2 or H-4), 8.51 (s, 1H, H-2 or H-4), 11.20 (br. s, 1H, NH). MS, *m/z* (%): 301 (M⁺, 49), 211 (32), 181 (42), 153 (57), 126 (94), 74 (100).

- 3 S. G. Zlotin, P. G. Kislitsin, A. I. Podgursky, A. V. Samet, V. V. Semenov, A. C. Buchanan III and A. A. Gakh, *J. Org. Chem.*, 2000, **65**, 8439.
- 4 I. L. Dalinger, T. I. Cherkasova, V. M. Khutoretskii and S. A. Shevelev, *Mendeleev Commun.*, 2000, 72.
- 5 X. Ouyang, N. Tamayo and A. S. Kiselyov, *Tetrahedron*, 1999, **55**, 2827.
- 6 C. Okafor, *Heterocycles*, 1977, **7**, 391.
- 7 K. Nagarajan, J. David, C. Kaul, R. Maller, R. Rao and R. Grewal, *Indian J. Physiol. Pharmacol.*, 1975, **19**, 39 (*Chem. Abstr.*, 1975, **83**, 172257u).

Received: 17th January 2001; Com. 01/1752