

Chemoenzymatic synthesis of optically active phosphinic analogues of S-substituted sulfur-containing amino acids

Kirill V. Alferov,^a Yurii N. Zhukov,^a Nikolai G. Faleev,^b Elena N. Khurs^a and Ralii M. Khomutov^{*a}

^a V. A. Engelhardt Institute of Molecular Biology, Russian Academy of Sciences, 119991 Moscow, Russian Federation. Fax.: +7 095 135 1405; e-mail: khomutov@genome.eimb.relarn.ru

^b A. N. Nesmeyanov Institute of Organoelement Compounds, Russian Academy of Sciences, 119991 Moscow, Russian Federation. Fax.: +7 095 135 5085; e-mail: ngfal@ineos.ac.ru

10.1070/MC2003v013n03ABEH001770

The interaction of racemic 1-amino-3-(methylthio)propylphosphinic acid with benzylthiol catalysed by pyridoxal-5'-phosphate-dependent L-methionine- γ -lyase affords (*R*)-1-amino-3-(benzylthio)propylphosphinic acid, which was converted into the (*R*)-isomers of phosphinic analogues of homocysteine and methionine.

1-Aminoalkylphosphinic acids, analogues of natural α -amino acids, display pronounced biological activity and are extensively applied in enzymological studies.^{1–3} However, only a few optically active compounds of this kind have been obtained until now. Thus, (*R*)-isomers of phosphinic analogues of alanine and phenylalanine have previously been prepared by the stereoselective alkylation of chiral Schiff bases derived from (1*R*,2*R*,5*R*)-2-hydroxypinan-3-one and the ethyl esters of aminomethyl-(diethoxymethyl)phosphinic acid by methyl iodide and benzyl chloride, respectively, while the stereoselectivity of the reaction strongly depended on the structure of the alkyl halide.⁴ The (*S*)- and (*R*)-isomers of 1-aminoalkylphosphinic acids, including (*S*)- and (*R*)-1-amino-3-(methylthio)propylphosphinic acids (analogues of methionine), have been obtained by separating diastereomeric salts of corresponding N-benzyloxycarbonyl derivatives with (*R*)- or (*S*)-methylbenzylamines.⁵ The prospects of using enzymatic methods for the separation and/or synthesis of chiral phosphinic analogues of amino acids were first demonstrated by the example of interaction of the enantiomers of a tyrosine phosphinic analogue with PLP-dependent L-tyrosine-phenol-lyase.³ Thus, only (*R*)-isomer of the phosphinic acid, whose configuration corresponds to natural (*S*)-tyrosine, proved to be the substrate in the reaction of α,β -elimination. Accordingly, (*R*)-isomer of the tyrosine phosphinic analogue was the product of the reverse enzymatic reaction.


In this work, a new method is proposed for the synthesis of optically active phosphinic analogues of S-substituted sulfur-containing amino acids. It is based on the use of L-methionine- γ -lyase for the transformation of racemic 1-amino-3-methylthiopropylphosphinic acid **1**[†] into optically active 1-amino-3-(benzylthio)propylphosphinic acid **2** and the subsequent preparation of chiral 1-amino-3-mercaptopropylphosphinic acid **3** followed by its S-methylation to known (*R*)-1-amino-3-methylthiopropylphosphinic acid **4**.

PLP-dependent L-methionine- γ -lyase [L-methionine-methanethiol lyase (deaminating) E.C. 4.4.1.11] from *Citrobacter intermedius* catalyses decomposition of L-methionine to methanethiol, α -ketobutyrate and the ammonium ion and displays specificity with respect to substrate structures and types of chemical reactions.⁷

An important feature of the enzyme is its ability to catalyse the enantioselective formation of S-alkylhomocysteins from L-methionine and thiols due to γ -substitution reaction. This reaction was also possible for analogue **1**, whose interaction with benzylthiol under the action of *Citrobacter intermedius* cells containing L-methionine- γ -lyase afforded optically active acid **2**, although with a low yield.⁸

We examined various conditions of the enzymatic synthesis of compound **2** from acid **1** and benzylthiol and found that the use of L-methionine- γ -lyase[‡] instead of intact cells as a biocatalyst allowed us to increase the yield of **2** up to 45%.⁸ The (*R*)-configuration was ascribed to optically active acid **2** based

[†] Racemic 1-amino-3-methylthiopropylphosphinic acid **1** was synthesised according to a published procedure.⁶



Scheme 1

on that the stereochemistry of the enzymatic reaction remains the same on going from the natural substrate to its phosphinic analogue, as it was observed for the reaction of L-tyrosine-phenol-lyase with the enantiomers of tyrosine phosphinic analogues. This assumption was supported experimentally by obtaining previously described phosphinic analogue **4** as a result of chemical transformation of amino acid **2** not affecting the chiral centre of the molecule. The debenzoylation of amino-phosphinate **2** by treatment with Na in liquid ammonia affords optically active amino acid **3**, which was not described before and may be used as a starting material for the synthesis of phosphinic analogues of various chiral S-substituted sulfur-containing amino acids.[†] By the methylation of analogue **3**, optically active acid **4** was prepared, for which the sign and value of specific rotation corresponded to (*R*)-isomer of the methionine phosphinic analogue⁵ (Scheme 1).^{††}

Thus, the (*R*)-configuration corresponds to amino acids **2–4**, which agrees with our assumption that the enantioselectivity of enzymatic γ -substitution reactions is retained and shows that phosphinic analogues of amino acids are stable with respect to racemisation under the action of Na in liquid ammonia.

[‡] L-Methionine- γ -lyase was obtained from *C. intermedius* cells according to a published procedure.⁸ The activity of the preparation was assayed by measuring the rate of α -ketobutyrate formation from L-methionine according to Friedemann.⁹ One unit of enzymic activity was determined as the enzyme amount catalysing the transformation of 1 μ mol of L-methionine per minute at 30 °C and a 40 mM concentration of L-methionine.

[§] Benzylthiol (0.5 ml) was added to a solution of **1** (169 mg, 1 mmol) in 20 ml of a 0.1 M potassium phosphate buffer containing 0.1 mM PLP and 12.8 U of L-methionine- γ -lyase. The reaction mixture was stirred on a shaker for three days at 30 °C. The protein was denatured by adding 30% trichloroacetic acid (2 ml) and removed by centrifugation. The solvent was evaporated *in vacuo*; the residue was dissolved in water (1 ml) and applied to a 40 ml column with Dowex 50 \times 8 resin (H⁺ form). The column was washed with water (100 ml), and product **2** was eluted with a 5% ammonia solution. The fractions containing **2** were evaporated *in vacuo* to dryness. The residue was dried *in vacuo* over P₂O₅ to give phosphinic analogue **2** (110 mg, 45%), mp 221 °C, $[\alpha]_D^{20}$ –16.3° (c 0.5, 1M HCl). *R*_f 0.61 (Pr⁺OH–25% NH₄OH–H₂O, 7:1:2), *R*_f 0.46 (BuOH–AcOH–H₂O, 12:3:5). ¹H NMR (400 MHz, 0.25 M NaOD in D₂O) δ : 1.63–2.05 (m, 2H, CH₂CH), 2.53–2.79 (m, 3H, SCH₂CH₂ and CH), 3.80 (s, 2H, CH₂Ph), 6.72 (dd, 1H, PH, *J* 486 Hz, *J* 1.8 Hz), 7.40 (s, 5H, Ph).

This work was supported by the Russian Foundation for Basic Research (grant nos. 03-04-48949 and 01-04-48636) and grant ‘Scientific school’ no. 1800-2003.4.

References

- 1 J. G. Dingwall, *Proceedings of the III Int. Conf. Chem. and Biotech. Biol. Active Comps.*, Sofia, 1985, vol. 1, p. 87.
 - 2 R. M. Khomutov, Yu. N. Zhukov, A. R. Khomutov, E. N. Khurs, D. L. Kramer, J. T. Miller and K. V. Porter, *Bioorg. Khim.*, 2000, **26**, 718 (*Russ. J. Bioorg. Chem.*, 2000, **26**, 647).
 - 3 N. G. Faleev, Y. N. Zhukov, E. N. Khurs, O. I. Gogoleva, M. V. Barbolina, N. P. Bazhulina, V. M. Belikov, T. V. Demidkina and R. M. Khomutov, *Eur. J. Biochem.*, 2000, **267**, 6897.
 - 4 P. P. McCleery and B. Tuck, *J. Chem. Soc., Perkin Trans. 1*, 1989, 1319.
 - 5 E. K. Baylis, C. D. Campbell and I. G. Dingwall, *J. Chem. Soc., Perkin Trans. 1*, 1984, 2845.
 - 6 T. I. Osipova, A. R. Khomutov, Yu. N. Zhukov and R. M. Khomutov, *Izv. Akad. Nauk, Ser. Khim.*, 1999, 1360 (*Russ. Chem. Bull.*, 1999, **48**, 1348).
 - 7 H. Tanaka, N. Esaki and K. Soda, *Biochemistry*, 1977, **16**, 100.
 - 8 K. V. Alferov, N. G. Faleev, E. N. Khurs, Y. N. Zhukov and R. M. Khomutov, *Mendeleev Commun.*, 2002, 2.
 - 9 F. Friedemann and G. Haugen, *Z. Biol. Chem.*, 1943, **177**, 415.
- † Sodium metal (20 mg, 0.86 mmol) was added to a solution of compound **2** (100 mg, 0.4 mmol) in 15 ml of boiling liquid ammonia. The reaction mixture was stirred for 30 min, and then solid NH₄Cl was added until the disappearance of a blue colour. The ammonia was allowed to evaporate, and the residue was concentrated with H₂O *in vacuo*. Thiol **3** was isolated on a 10 ml column with Dowex 50×8 resin (H⁺ form) by elution with a 15% aqueous isopropanol solution. The fractions containing **3** were evaporated *in vacuo* to dryness. The residue was dried *in vacuo* over P₂O₅ to give phosphinic analogue **3** (27 mg, 43%). [α]_D²⁰ –22° (c 1, H₂O). R_f 0.39 (PrⁱOH–25% NH₄OH–H₂O, 7:1:2), R_f 0.23 (BuOH–AcOH–H₂O, 12:3:5). ¹H NMR (400 MHz, D₂O) δ: 1.80–2.06 (m, 2H, CH₂CH), 2.48–2.78 (m, 3H, SCH₂), 3.21–3.25 (m, 1H, CH), 6.87 (d, 1H, PH, *J* 535 Hz).
- †† MeOH (1.4 ml) and MeI (0.01 ml) were added to a solution of compound **3** (25 mg, 0.15 mmol) in 0.15 ml of 2 M NaOH. The mixture was allowed to stand for 16 h and concentrated *in vacuo* to dryness. Compound **4** (15 mg, 59%) was isolated as described above. Mp 230 °C, [α]_D²⁰ –31° (c 1, H₂O) {lit.,⁵ [α]_D²² –30° (c 1, H₂O)}. R_f 0.53 (PrⁱOH–25% NH₄OH–H₂O, 7:1:2), R_f 0.36 (BuOH–AcOH–H₂O, 12:3:5). ¹H NMR (400 MHz, D₂O) δ: 1.8–2.28 (m, 2H, CH₂CH), 1.97 (s, 3H, MeS), 2.51–2.65 (m, 2H, CH₂S), 3.14–3.19 (m, 1H, CH), 6.87 (d, 1H, PH, *J* 535 Hz).

Received: 21st April 2003; Com. 03/2096