
Unexpected reaction of chitosan with 2-ethoxy-1-ethoxycarbonyl-1,2-dihydroquinoline

Evgeniya A. Stepnova, Vladimir E. Tikhonov,* Tatyana A. Babushkina, Zinaida S. Klemenkova and Igor A. Yamskov

A. N. Nesmeyanov Institute of Organoelement Compounds, Russian Academy of Sciences, 119991 Moscow, Russian Federation.
Fax: +7 495 135 5085; e-mail: tikhon@ineos.ac.ru

DOI: 10.1070/MC2006v016n05ABEH002363

The direct reaction of 2-ethoxy-1-ethoxycarbonyl-1,2-dihydroquinoline (EEDQ) with chitosan amino groups, which resulted in the formation of *N*-(ethoxycarbonyl)chitosan, was found and confirmed chemically and by IR and ¹H NMR spectroscopy.

Chitosan is a partially or fully *N*-deacetylated natural polysaccharide chitin and is considered to be a copolymer of *D*-glucosamine and *N*-acetyl-*D*-glucosamine. Chitosan is of considerable interest since it is a biodegradable, nontoxic cationic polymer. During the last decades, chitosan has found numerous applications in medicine, cosmetics and food technology.¹ Chitosan has the apparent $pK_a \sim 6.5$ and is only soluble in acidic aqueous solutions with $pH < 6.0$, and under these conditions chitosan can react with carboxylic acids in the presence of a coupling reagent.² A widely used coupling reagent is 2-ethoxy-1-ethoxycarbonyl-

1,2-dihydroquinoline (EEDQ).³ EEDQ is a stable, readily available reagent, which has been used for smooth preparation of amides and peptides in high yields,^{4–6} as well as in neurochemical and pharmaceutical researches^{7,8} and affinity sorbent preparations.⁹ In organic solvents, EEDQ has been used for protecting the hydroxyl groups of 2-hydroxycarboxylic acids since the *O*-ethoxycarbonyl groups can be easily removed under weakly basic conditions.⁴ For these reasons, EEDQ is a very attractive coupling reagent, especially because it fails to react with amines under conditions of peptide synthesis.^{6,10}

- 5 B. Zacharie, T. P. Connolly and C. L. Penney, *J. Org. Chem.*, 1995, **60**, 7072.
- 6 B. Belleau and G. Malek, *J. Am. Chem. Soc.*, 1968, **90**, 1651.
- 7 K. Y. Vinod, M. N. Subhash and B. N. Srinivas, *Neurochem. Res.*, 2001, **26**, 113.
- 8 B. Floran, B. Gonzales, L. Floran, D. Erlij and J. Aceves, *Eur. J. Pharm.*, 2005, **520**, 43.
- 9 M. H. Hyun, J. S. Na and C.-S. Min, *J. Chromatogr. A*, 1996, **732**, 209.
- 10 B. Belleau, R. Martel, G. Lacasse, M. Menard, N. L. Weinberg and Y. G. Perron, *J. Am. Chem. Soc.*, 1968, **90**, 823.
- 11 A. Hirai, H. Odani and A. Nakajima, *Polym. Bull.*, 1991, **26**, 87.
- 12 J. Kaczaj, *Appl. Spectrosc.*, 1967, **21**, 180.
- 13 D. Ben-Ishai and A. Berger, *J. Org. Chem.*, 1952, **17**, 1564.
- 14 *Beilstein Handbuch Der Organische Chemie*, ed. F. Richter, Verlag von Julius Springer, 1929, Auflage 4, Band 3/4, S. 10.
- 15 C. Muzzarelli, G. Tosi, O. Francescangeli and R. A. A. Muzzarelli, *Carbohydr. Res.*, 2003, **338**, 2247.
- 16 S. L. Johnson and D. L. Morrison, *J. Am. Chem. Soc.*, 1972, **94**, 1323.

Received: 29th March 2006; Com. 06/2708